

Identifying essential COVID-19 indicators for primary healthcare through Delphi analysis in 31 European countries: Eurodata eDelphi study

Maria Pilar Astier Peña^{1,2,3,4,+}, Raquel Gómez Bravo^{5,6,7,+}, Ileana Gefaell Larrondo⁸, Lourdes Ramos Del Rio⁹, José Joaquín Mira¹⁰, Snežana Knežević¹¹, Aleksandar Kirkovski¹², Büsra Çimen Korkmaz¹³, Milena Kostić¹⁴, Anna Krztoń-Królewiecka¹⁵, Anna Segernäs¹⁶, Heidrun Lingner¹⁷, Liubovė Murauskienė¹⁸, Achim Mortsiefer¹⁹, Katarzyna Nessler²⁰, Nagu Penakacherla²¹, Maria Pencheri²², Ábel Perjés²³, Ferdinando Petrazzuoli²⁴, Goranka Petricek²⁵, Theresa Sentker¹⁷, Lucia Palandri²⁶, Davorina Petek²⁷, Bert Vaes²⁸, Oksana Ilkov²⁹, Erva Kırkoç Üçüncü³⁰, Shlomo Vinker³¹, Radost Assenova³², Limor Adler³¹, Maria Bakola³³, Sherihane Bensemmane³⁴, Ludmila Bezdícková³⁵, Sabine Bayen³⁶, Jako S. Burgers³⁷, Carmen Busneag³⁸, Georgi Tsigarovski³², Asja Cosic Divjak³⁹, Philippe-Richard J. Domeyer⁴⁰, Louise Fitzgerald^{41,42}, Dragan Gjorgjievski⁴³, Bruno Heleno^{44,45}, Kathryn Hoffmann⁴⁶, Marijana Jandrić-Kocić⁴⁷, Ana Luísa Neves^{48,49}, Marina Guisado-Clavero⁵⁰, Sara Ares-Blanco^{9,51,52,*,*}, Thomas Frese^{53,*} and Eurodata Collaborative Group^{**}

¹Universitas Health Centre, Zaragoza, Spain

²Health Service of Aragon, GILS011, Instituto de Investigación Sanitaria de Aragón, Zaragoza, Spain

³QISP-Tar Research Group, Fundació Institut Universitari per a la Recerca a l'Atenció Primària de Salut—IDIAP Jordi Gol, Barcelona, Spain

⁴Aragones Health Service, Patient Safety Working Party of semFYC (Spanish Society for Family and Community Medicine), Quality and Safety in Family Medicine of WONCA World (Global Family Doctors), Board Member of WONCA World, Barcelona, Spain

⁵CHNP, Rehaklinik, Ettelbruck, Luxembourg

⁶Research Group Self-Regulation and Health, Institute for Health and Behaviour, Luxembourg, Luxembourg

⁷Department of Behavioural and Cognitive Sciences, Faculty of Humanities, Education, and Social Sciences, Luxembourg University, Esch-sur-Alzette, Luxembourg

⁸Fundación de Investigación e Innovación Biosanitaria de Atención Primaria (FIIBAP), Research Network on Chronicity, Primary Care and Health Promotion-RICAPPS (RICORS), Spain

⁹Federica Montseny Health Centre, Gerencia Asistencial de Atención Primaria, Servicio Madrileño de Salud, Madrid, Spain

¹⁰Universidad Miguel Hernández, Elche, Spain

¹¹Department of Medical Sciences, Academy of Applied Studies Polytechnic, Belgrade, Serbia

¹²Faculty of Medicine, Ss. Cyril and Methodius University, Skopje, North Macedonia

¹³Homecare Unit, Van Gürpınar District Hospital, Gürpınar, Van, Turkey

¹⁴Health Center “Dr Đorđe Kovačević”, Lazarevac, Belgrade, Serbia

¹⁵Department of Family Medicine, Andrzej Frycz Modrzewski Krakow University, Krakow, Poland

¹⁶Primary Health Care Centre in Ekholmen and Department of Health, Medicine and Caring Sciences, Linköping University, Linköping, Sweden

¹⁷Department of Medical Psychology OE 5430, Hannover Medical School, Hanover, Germany

¹⁸Department of Public Health, Institute of Health Sciences, Faculty of Medicine, Vilnius University, Vilnius, Lithuania

¹⁹Chair of General Practice II and Patient-Centredness in Primary Care, Institute of General Practice and Primary Care, Faculty of Health, Witten/Herdecke University, Witten, Germany

²⁰Department of Family Medicine, Uniwersytet Jagielloński—Collegium Medicum, Kraków, Poland

²¹Portfolio GP, Handsworth Medical Practice, Birmingham, UK

²²Department of Public Health, Institute of General Medical Practice, Aarhus University, Aarhus, Denmark

²³Department of Family Medicine, Semmelweis University, Budapest, Hungary

²⁴Department of Clinical Sciences in Malmö, Centre for Primary Health Care Research, Lund University, Malmö, Sweden

²⁵Department of Family Medicine “Andrija Stampar” School of Public Health, School of Medicine, University of Zagreb, Health Centre Zagreb West, Zagreb, Croatia

²⁶Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy

²⁷Department of Family Medicine, Faculty of Medicine, University of Ljubljana, Ljubljana, Slovenia

²⁸Department of Public Health and Primary Care, KU Leuven, Leuven, Belgium

²⁹Department of Family Medicine and Outpatient Care, Medical Faculty 2, Uzhhorod National University, Uzhhorod, Ukraine

³⁰Home Health Services Unit, Egil District State Hospital, Diyarbakir, Turkey

³¹Department of Family Medicine, Faculty of Medical & Health Sciences, Tel Aviv University, Tel Aviv, Israel

³²Department of Urology and General Practice, Faculty of Medicine, Medical University of Plovdiv, Plovdiv, Bulgaria

³³Research Unit for General Medicine and Primary Health Care, Faculty of Medicine, School of Health Science, University of Ioannina, Ioannina, Greece

³⁴Department of Epidemiology and Public Health, Health Services Research, Sciensano, Brussels, Belgium

³⁵Department of Postgraduate Medical Education and Czech Society of General Practice, Prague, Czech Republic

³⁶Department of General Practice, University of Lille, UFR3S, Lille, France

³⁷Department of Family Medicine, Care and Public Health Research Institute, Maastricht University, Maastricht, The Netherlands

³⁸“Spiru Haret” University, Bucharest, Romania

³⁹Health Centre Zagreb Centar, Zagreb, Croatia

⁴⁰School of Social Sciences, Hellenic Open University, Patra, Greece

⁴¹Member of Irish College of General Practice (MICGP), Dublin, Ireland

⁴²Member of Royal College of Physician (MRCSI), Dublin, Ireland

⁴³Center for Family Medicine, Medical Faculty Skopje, Skopje, North Macedonia

⁴⁴Comprehensive Health Research Center, NOVA Medical School, Universidade Nova de Lisboa, Lisbon, Portugal

⁴⁵USF das Conchas, Regional Health Administration Lisbon and Tagus Valley, Lisbon, Portugal

⁴⁶Department of Primary Care Medicine, Medical University of Vienna, Vienna, Austria

⁴⁷Health Center Krupa na Uni, Bosnia and Herzegovina, Republic of Srpska

⁴⁸Department of Primary Care and Public Health, Imperial College London, London, UK

⁴⁹CINTESIS@RISE, Faculty of Medicine of the University of Porto, Porto, Portugal

⁵⁰Investigation Support Multidisciplinary Unit for Primary Care and Community North Area of Madrid, Gerencia Asistencial Atención Primaria, Servicio Madrileño de Salud, Madrid, Spain

⁵¹Research Network on Chronicity, Primary Care and Health Promotion-RICAPPS (RICORS), Spain

⁵²Instituto de Investigación Sanitaria Gregorio Marañón, Madrid, Spain

⁵³Institute of General Practice and Family Medicine, Martin-Luther University Halle-Wittenberg, Halle/Saale, Germany

Corresponding author. Centro de Salud Federica Montseny, Avenida Albufera, 285, 28038 Madrid, Spain.

E-mail: sara.ares@salud.madrid.org.

†These authors contributed equally to the first and second position of this work.

‡These authors contributed equally to the senior position of this work.

**Members of the Eurodata collaborative groups are listed in the Acknowledgements section.

Abstract

Background: The COVID-19 pandemic has underlined the essential role of primary healthcare (PHC) in epidemiological surveillance and public health decision-making. Across Europe, the integration of electronic health records (EHRs) and the sentinel networks have been pivotal in monitoring COVID-19. However, the lack of standardized PHC indicators for COVID-19 hinders the comparability of data among countries. **Objective:** To establish a consensus on a set of standardized PHC activity indicators related to the COVID-19 pandemic for 31 countries, enhancing the capability of health authorities to make informed decisions and prepare for future health crises. **Methods:** A two-round eDelphi study was conducted using a structured web-based survey, following the CREDES guidelines, to achieve consensus among a panel of 164 experts from the Eurodata study. 86 Indicators were selected based on their availability during the current pandemic, with participants rating the relevance and utility of proposed indicators. **Results:** Of the 22 initial indicators, seven received consensus for inclusion, while two remained contentious after the second round. The study found significant discrepancies in the awareness of sentinel networks and accessibility to PHC data. The consensus emphasized the necessity for indicators to be standardized, reproducible, and easily extractable from databases, with recommendations for disaggregation by age, sex, and vaccination status. **Conclusion:** Key COVID-19 indicators for PHC were identified, reflecting a consensus among healthcare professionals. Further cooperation between PHC providers and national public health authorities is warranted both on the national and the international level to harmonized healthcare indicators in response to future health emergencies.

Introduction

Epidemiological surveillance serves not only to detect cases and quantify disease trends but also as a foundation for health policy decision-making to face health crisis [1]. The COVID-19 pandemic underscored the importance of current robust surveillance systems, for COVID-19 and also for influenza and other respiratory viruses. The European Centre for Disease Prevention and Control (ECDC) and the World Health Organization (WHO) have emphasized the importance of gathering data from both primary and secondary healthcare settings. This approach is pivotal for detecting and responding to increases and outbreaks of acute respiratory infections throughout Europe [2].

The majority of European countries have adopted and integrated electronic health records (EHRs) in PHC. These EHRs played a relevant role as valuable sources of data for both collection and monitoring of the COVID-19 pandemic's progression in some European countries [3], especially since the majority of COVID-19 patients received outpatient care in PHC [4]. They provided insights into the volume of acute respiratory infection (ARI) consultations [5], the healthcare worker attending patients [6], sick leave statistics [7, 8], and prescribed treatments. At the same time, the role of primary care sentinel networks has been well-established. Originally focused on detecting influenza-like illness (ILI), the definition has now

expanded to include ARI, enabling the collection of data on COVID-19 and other viral infections [9, 10]. Nowadays, 29 European countries are reporting influenza data of ILI and ARI consultations. Currently, the general practitioners (GPs) participating in these sentinel networks represent 1-5% of the total number of GPs in their countries [9]. Despite their importance, support for these networks is inconsistent; some countries, like the Czech Republic, operate on voluntary participation without funding. The coexistence of sentinel networks and EHRs in Europe has marked a new era for monitoring COVID-19 in PHC [11]. Many countries have leveraged both sources to report the pandemic's progression within their communities, resulting in the creation of over 40 distinct COVID-19 PHC indicators across Europe [4]. At an European level, the ECDC is collecting data based from primary care mainly from the sentinel networks [12], public European data concerning other PHC COVID-19 indicators remains scarce. It is important to remember the lack of a common application programming interface, which limits the interoperability of EHRs, as well as the infrequent use of open data standards. The objective of this study is to establish a consensus on a set of PHC indicators that describe the workload in PHC related to diagnosing, follow-up, and responding to patients' needs during the COVID-19 pandemic across 31 European countries through a web-based Delphi study [13]. The overarching goal is to equip European health authorities with a cohesive PHC scorecard,

thus enhancing their capacity to make informed policy decisions regarding organizing work in health centers, allocating resources, and assessing PHC team in a collaborative response within national health systems to future health crises across the continent.

Methods

Study design

The Delphi technique employs a multistage self-administered questionnaire with individual feedback, to ascertain consensus from a larger group of experts [14, 15]. To ensure methodological rigor, we used the CREDES guideline [16], available in [Supplement S1](#), along with the invitation letter included in [Supplement S2](#).

Development of eDelphi survey

We selected PHC indicators available in 31 European countries during the ongoing COVID-19 pandemic identified in a previous Eurodata project study [4]. In addition to this, a comprehensive literature review was performed to enrich the list. Following thorough discussions within the research team, a definitive list of indicators was selected for inclusion in the eDelphi survey. This survey consisted of 85 items, organized into 11 distinct sections as detailed in [Supplement S5](#). Each eDelphi round consisted of an introductory part on the purpose of each round. To avoid confusion among the different levels of decision-making (practice, regional, national, or European level). The process of constructing the eDelphi survey is depicted in [Fig. 1A](#). Then, the research team discussed and decided on the final list of indicators to consider for the eDelphi survey.

An initial version of the survey was shared with the Eurodata research network, aiming to achieve a consensus on the included items [17], clarify the various decision-making levels, and define each indicator precisely. Subsequent to a review and integration of all the feedback, a revised draft was prepared. To ensure its validity, a pilot survey was conducted with a selection of Eurodata researchers invited to provide feedback. Incorporating this input, the final questionnaire was then configured on the online platform. Once the final questionnaire was designed, it was shared with the participants so they could respond.

We defined consensus as at least 70% of respondents agreeing or strongly agreeing on a given item within the team. Additionally, it was required that at least 40% of this 70% rate the item as “strongly agree” to achieve consensus [18]. Non-consensus was defined as less

than 70% of respondents agreed or strongly agree, and no major change was suggested for the item. Those indicators with less than 70% and more than 50% in consensus ratings were sent to the second round. Those with less than 50% in consensus ratings and without any suggestions for change by the panel after two rounds were rejected for the set of indicators.

Study setting and expert panel

We recruited 164 panelists from 31 European countries with prior involvement in or connections to the Eurodata project, allowing us to form a larger group in a cost-effective manner ([Fig. 1B](#)). Selection criteria were (i) to be a front-line family doctors involved in PHC COVID-19 pandemic, (ii) belonging to WONCA Europe Organization or EGPRN (European General Practitioner Research Network), and (iii) Public health technical officer linked to PHC in any European countries participating in the study and being proposed by a family doctor linked to the project.

Data collection and analysis

The online form was built up in the Delphi Studies Platform of Miguel Hernandez University of Elche (Spain). To collect our data, we used the Qualtrics XM Platform. A personal link was sent via email to each panelist. The researchers on the team did not have access to the responses, which were coded anonymously, preventing the extraction of stratified data that could break anonymity. This allowed following up response rates and sending reminders to specific members. Due to high workload caused by the COVID-19 pandemic, each round lasted 8 weeks. We opted for a flexible approach towards the panelists to increase the response rate of each round. Reminders were sent weekly sent to members that had not completed the survey. Data collection took place between October 2022 and March 2023. For analyzing quantitative data, we calculated descriptive statistics of every item using SPSS 27 (IBM SPSS Statistics 27). We used Microsoft Excel to list and categorize qualitative data. Panelists' comments were anonymously and literally registered. For analyzing qualitative data, we used content analysis.

Role of research team to prevent bias

The research team made methodological decisions in line with the available literature. We predefined and stipulated methodological steps before commencing the study. We applied, monitored, and evaluated these steps during the study. The results of each round were discussed





Figure 1. (A) Process steps for constructing the eDelphi survey. (B) Participants invited to answer the eDelphi and countries of origin.

by the research team, while qualitative data were interpreted by two researchers for researcher triangulation (Supplement S1).

Results

A two-wave eDelphi consensus survey has been performed. The first round of participation included 117 respondents (71.3%) between October and December 2022, while the second round saw 101

respondents (86.3%) between January and March 2023, despite two personalized email reminders sent during each round. Mean age was 44.1 years, with 69.1% males, 62.2% were GPs and 30.9% were public health officers (Table 1). Regarding years of work experience, 61% had more than 3 years of experience. Country of origin are provided in Fig. 1B and Supplement S3.

In questions regarding the conceptualization of health indicators, the majority of participants concurred on the importance of

Table 1. Demographic information of eDelphi participants and utility of health indicators in pandemic monitoring and data disaggregation levels (eDelphi first round, *n*: 117)

Demographic information of eDelphi participants						
Sex	<i>n</i>	Professional background (<i>n</i>)				
Male	84	General Practice			76	
Female	32	Public Health			38	
Unknown	1	Other			8	
Age groups		Unknown			5	
<35 years	23	Years of expertise (<i>n</i>)				
35–49 years	54	<5 years			28	
>50 years	36	5–9 years			16	
Unknown	4	10–19 years			40	
		20–29 years			21	
		>30 years			12	
Utility of health indicators in pandemic monitoring and data disaggregation levels						
Level of agreement on values relevant for a health indicator and level of disaggregation		Strongly disagree (%)	Disagree (%)	Neither agree nor disagree (%)	Agree (%)	Strongly agree (%)
Values considered relevant to create an indicator in primary care						
• Health indicators attempt to describe and monitor a population's health status or condition		1.7	0.8	3.4	31.9	62.2
• An indicator is a measurement that reflects health characteristics in a given population		0.8	4.2	11.8	35.3	47.9
• Indicators are dynamic, reflecting a specific time-linked period		1.7	3.4	10.1	38.7	46.2
• Health indicators can be used to describe disease burden in a specific population group		0.8	1.7	12.6	34.5	50.4
• Health indicators can be used to forecast the risk of disease outbreaks and helping to prevent epidemic/pandemic		2.6	5.1	18.8	37.6	35.9
• Indicators are used in public health to drive decision-making for the health of the community		1.7	3.5	8.7	33.0	53.0
• Regular monitoring indicators can provide feedback to improve decision-making in healthcare systems		0.9	0.0	6.1	34.8	58.3
• Health indicators should have a common and clear definition for all primary care providers		0.9	1.7	3.5	18.3	75.7
• Health indicators should be reproducible		0.0	0.0	2.6	23.5	73.9
• Health indicators should be feasible and designed to allow for easy extraction from database		0.0	1.8	5.3	23.7	69.3
Population disaggregation level of indicators						
• COVID-19 primary care indicators should be disaggregated by sex		0.8	9.2	16.8	16.8	56.3
• COVID-19 primary care indicators should be disaggregated by age		0.0	0.8	4.2	12.6	82.4
• COVID-19 primary care indicators should be disaggregated by vaccination status		0.0	2.5	6.8	19.5	71.2
• COVID-19 primary care indicators should be disaggregated by group ethnic and/or migrant situation		5.9	10.9	22.7	25.2	35.3
• COVID-19 primary care indicators should be disaggregated by vulnerable populations (low socioeconomic status, health illiteracy, homeless people, etc.)		0.8	5.0	10.9	22.7	60.5
• Would you like to add other indicator regarding this section?		Urban/rural; Local/Regional; Comorbidities; Functional status; war; PHC system (free of charge or co-payment)				

Bold type denotes $\geq 70\%$ of agreement throughout participants.

indicators and the necessity for a common definition (Table 1). A majority of participants, 71.6%, were not aware of a sentinel doctor network in their countries. Additionally, 67.5% reported a lack of public PHC data availability, while 57.9% indicated access to PHC data from insurance companies. For 61% of the participants, PHC data were available at the practice level, encompassing both public and insurance company sources. However, only 40% had access to regional and national PHC data from both public and private providers.

Table 1 summarizes the results of the first and second round of Delphi survey with the accepted or rejected indicators. In the first round of the Delphi survey, four items were accepted and seven were rejected. Subsequently, 11 items advanced to the second round, where 3 were accepted, 6 rejected, and 2 remained unclear for acceptance. In terms of content, all items related to the sentinel network and suspected COVID-19 cases were accepted. However, items pertaining to nurse follow-up of patients, as well as those concerning home visits and sick leaves, were uniformly rejected.

Regarding the chosen level of data disaggregation, the majority of participants (80-96%) indicated having access to regional and national-level information. In terms of publication frequency, most respondents mentioned daily reporting during pandemic peaks and weekly reporting during other periods (96%). As for proposed new indicators, the suggestions primarily revolved around revising some from the survey. Suggestions such as gathering information during home visits, conducting follow-ups, documenting the types of symptoms observed when patients were referred to the hospital, or detailing the types of complementary tests conducted were shared. Comments and proposal of new indicators were recorded (Supplement S4 (S4-1 and S4-2)).

In terms of indicators for estimating suspicious COVID-19 cases, the majority of comments were focused on the definition of denominators and how to collect the data (Tables 1 and 2). In terms of follow-up, one noteworthy aspect was the number of COVID-19 patients who received follow-up care solely in PHC, without requiring hospitalization. The accepted and uncertain indicators are described in Table 3.

Discussion

Main findings

The results of this comprehensive eDelphi study provide a pivotal step towards the standardization of PHC indicators in the context of the COVID-19 pandemic across Europe. PHC data availability is scarce across countries participants. There is a need to define health indicators integrating PHC workload in order to tailor public health policies. Most countries adopted health measures during the COVID-19 pandemic based upon data collected from secondary care (hospital admissions being the most important measure in most countries), whereas data from PHC which took care of the majority of COVID-19 patients were not used as effectively. Seven indicators have been defined, considering COVID-19 estimated cases. Robust agreement on indicators related to case detection and management within the sentinel networks were found. These indicators are valid not only for the EHR systems but also for sentinel networks.

These proposed indicators for COVID-19 should enable us to develop robust indicators for surveillance of respiratory illness in PHC (not only during pandemics) and mainly connect those to the national public health authorities, which collect data for international surveillance. The robust agreement on indicators related to case detection and management within the sentinel networks underscores the critical role these networks play in national surveillance systems. These networks facilitate real-time data collection, which is vital for the timely response to infectious disease outbreaks [9, 10]. However, variations in the denominators used for these indicators highlight the challenges of harmonizing data across diverse

European healthcare systems. Although there is strong agreement on the usefulness of sentinel networks, not all GPs were aware of the existence of the sentinel network, highlighting the need to promote the network and make the results public. Sentinel networks must be valued and given sufficient support (both financial and technical); otherwise, the quality of the collected data will not be adequate to draw valid conclusions.

The number of face-to-face visits to GPs by COVID-19 patients and the count of physical examinations were selected as key indicators. These indicators are essential for assessing the burden of disease and the capacity of PHC systems to manage patient loads. However, indicators related to the frequency of home visits, and the role of nurses were rejected. These results highlight the varying contexts in which different health systems and PHC services operate, such as differences in patient access and the need for sick leave policies. European PHC systems showed variation, especially in nursing roles, that were not involved in clinical encounters with COVID-19 patients in countries such as Austria, Czech Republic, Germany, Greece, Italy, Luxemburg, Bulgaria, and Belgium [6]. However, they are in charge of long-term facilities and home visits in regular demand instead of GPs in many European countries in a daily basis [6, 19]. This finding warrants further investigation, as nursing services have been integral to the pandemic response, especially in community settings and for patients requiring long-term care [19]. Nevertheless, community settings and long-term care are not considered as part of PHC facilities services in many European countries. As PHC increasingly adopts multi-professional team structures, monitoring the activities of other professionals becomes pertinent, particularly during staffing shortages experienced at the peak of the pandemic.

Other indicators that were rejected pertained to sick leaves. The ethical dilemma of self-declared sick leave from self-testing, as France proceed, has emerged and it opens again the discussion of self-empowerment due to the fact that a self-declaration of a positive test provided an online sick leave [20]. As studies highlight a decrease in essential PHC activities during the pandemic [21–24], there is a pressing need to enhance legislation across Europe. This is crucial to encourage self-declared sick leave for mild COVID-19 cases, preventing the overburdening of PHC with low-priority tasks and promoting high-value care activities [25].

Also, the burden of administrative work provided by PHC has become no longer bearable and unwarranted with regard to uncomplicated patients in whom a sick-leave could be managed without the need for a GP consultation.

It also notes the importance of tracking hospital referred patients (percentage of COVID-19 patients treated who are admitted to the hospital). The referred activity from hospital to PHC could be defined as the percentage of COVID-19 patients treated who are discharged [COVID-19 rate (cases and/or contacts) per 10 000 assigned inhabitants].

Indicators related with complementary test have been evaluated too. The study suggests separating X-ray and blood test demands indicators and stratifying by COVID-19 severity. It might offer a proxy of the complexity of COVID-19 patients treated in PHC. Nevertheless, there is a high variability on the emergency X-ray access for GPs among European countries [6].

The inclusion of mandatory reporting of COVID-19 cases in PHC as an accepted indicator in the initial round holds significance as it marked the foundational step toward constructing a PHC scorecard related to COVID-19 activity, interconnected with public health information. The ease of collecting this indicator stemmed from certain countries' readiness to integrate their PHC Information Systems with PHC EHR and public health information systems, allowing for electronic mandatory disease reporting [26]. Including this indicator as the primary measurement to monitor the pandemic could be an excellent initial step in a comprehensive scorecard for tracking the pandemic [27, 28]. This could complement metrics like the number of COVID-19 hospitalizations or ICU

Table 2. Round 1 and round 2 from the consensus of eDelphi items with mean scores

	Results Round: 1						Results Round: 2					
	N	Mean	VC (%)	%>=4	%=5	Decision	N	Mean	VC (%)	%>=4	%=5	Decision
Indicators regarding role sentinel network												
• Positive cases for SARS-CoV-2 (COVID-19) seen by the Sentinel networks. Definition: Numerator: Positive cases for SARS-CoV-2 (COVID-19) seen by the Sentinel networks. Denominator: Total population of a country or region	116	4.1	22.8	73.3	42.2	Accepted						
• Positivity rates to SARS-CoV-2 (COVID-19) among all the respiratory infections by the Sentinel networks. Definition: Numerator: Positivity rates to SARS-CoV-2 (COVID-19) among all the respiratory infections by the Sentinel networks. Denominator: Total population of a country or region	115	4.1	22.8	80.9	40.0	Accepted						
• Estimated incidence of COVID-19 cases per 100 000 population with respiratory signs observed in general practice through the Sentinel networks. Definition: Numerator: Number of COVID-19 cases with respiratory signs from Sentinel network. Denominator: Region or country's total population	116	4.1	24.6	76.7	39.7	Second Round	102	4.1	20.3	74.51	39.2	Accepted
Indicators regarding suspicious COVID-19 cases												
• Percentage of cases of COVID-19 among all respiratory infection cases in PHC. Definition: Numerator: Number of cases of COVID-19 in primary care, region or country. Denominator: Number of all respiratory infection cases in primary care, region or country	116	4.1	22.1	78.4	37.9	Second Round	100	4.2	21.5	86	44	Accepted
• Total COVID-19 cases with positive test in primary care. Definition: Numerator: Number of primary care COVID-19 cases with positive test in practice, region or country. Denominator: Total primary care COVID-19 Tests performed in practice, region or country	117	4.1	24.3	75.2	40.2	Accepted						
Indicators regarding primary care follow-up to COVID-19 patients												
• Number of COVID-19 patients who were followed-up in primary care (nurse and/or GP) for all reasons. Definition: Numerator: Number of COVID-19 patients who were followed-up in primary care in a period. Denominator: Total number of patients visited in primary care in a period	117	3.9	28.8	70.1	38.5	Second Round	100	3.8	28.6	65	35	Doubtful
• Primary care follow-up ratio (nurses and/or GP): follow-up ratio of cases and contacts. Definition: Numerator: Family medicine follow-up COVID-19 cases. Denominator: Family medicine follow-up COVID-19 contacts	117	3.4	39	54.7	24.8	Second Round	101	3.2	42.1	46.53	20.8	Rejected
Indicators regarding the follow-up of primary care nurses to COVID-19 patients												
• Number of any contacts with nurse with COVID-19 recorded as reason for the contact. Definition: Numerator: Number of any contacts with nurse with COVID-19 recorded as reason for the contact. Denominator: Total contacts with nurse in a period	116	3.3	38.7	49.1	18.1	Rejected						
• Number of nurse home visits with COVID-19 recorded as reason for home care. Definition: Numerator: Number of nurse home visits with COVID-19 recorded as reason for home care. Denominator: Total nurse home visits with nurse in a period	115	3.3	38.8	47.8	20.0	Rejected						

(continued)

Table 2. Continued

	Results Round: 1						Results Round: 2					
	N	Mean	VC (%)	%>=4	%=5	Decision	N	Mean	VC (%)	%>=4	%=5	Decision
<ul style="list-style-type: none"> • Number of nurse telephone consultations with COVID-19 recorded as the reason for consultation. Definition: Numerator: Number of nurse telephone consultations with COVID-19 recorded as the reason for consultation. Denominator: Total telephone contacts with nurse in a period 	115	3.2	40.8	47	17.4	Rejected						
Number of nurse control home visit with COVID-19 recorded as the reason for home visit	113	2.9	46.5	37.2	14.1	Rejected						
Indicators regarding the follow-up of COVID-19 patients in primary care												
<ul style="list-style-type: none"> • Number of phone consultations to patients with COVID-19 or patients close family member (by GP). Definition: Numerator: Number of phone consultations to patients with COVID-19 or patients close family member by GP. Denominator: Total number of phone consultations to patients by GP 	117	3.6	33.7	60.7	27.4	Second Round	101	3.5	33.6	52.48	20.8	Rejected
<ul style="list-style-type: none"> • Number of email consultations to patients with COVID-19 or patients close family member (by GP). Definition: Numerator: Number of email consultations to patients with COVID-19 or patients close family member (by physician). Denominator: Total number of email consultations to patients by GP 	117	3.1	42.6	41	17	Rejected						
<ul style="list-style-type: none"> • Number of face-to-face visits to GP with COVID-19 recorded as reason for the visit. Definition: Numerator: Number of face-to-face visits to GP with COVID-19 recorded as reason for the visit. Denominator: Total number of face-to-face consultations to patients by GP 	117	4.0	25.2	72.6	40.2	Accepted						
<ul style="list-style-type: none"> • Number of first visits (examinations) with COVID-19 recorded as reason for the visit (by GP) 	113	3.8	27.1	65.5	31.9	Second Round	101	3.9	25.2	74.26	28.7	Accepted
<ul style="list-style-type: none"> • Number of control visits (examinations) with COVID-19 recorded as reason for the visit (by GP). Definition: Numerator: Number of first visits (examinations) with COVID-19 recorded as reason for the visit (by GP). Denominator: Total number of first visits to patients by GP 	114	3.5	34.2	59.6	23.7	Second Round	97	3.5	31.5	56.7	18.6	Rejected
<ul style="list-style-type: none"> • Number of first home visits with COVID-19 recorded as reason for home visit (by GP). Definition: Numerator: Number of first home visits with COVID-19 recorded as reason for the home visit (by GP). Denominator: Total number of first home visits to patients by GP 	113	3.6	32.7	60.2	25.7	Second Round	96	3.4	33.4	53.13	20.8	Rejected
<ul style="list-style-type: none"> • Number of follow-up home visits with COVID-19 recorded as reason for home visit (by GP). Definition: Numerator: Number of control home visits with COVID-19 recorded as reason for home visit (by physician). Denominator: Total number of control home visits to patients by GP 	114	3.3	37.1	48.2	21.1	Rejected						
Indicators regarding number of procedures in PHC to COVID-19 patients												
<ul style="list-style-type: none"> • Total number of procedures to patients in primary care with COVID-19 recorded as reason for procedures. Definition: Numerator: Total number of procedures to patients in primary care with COVID-19 recorded as reason for procedures. Denominator: Total number of procedures to all patients in the practice in a period 	116	3.4	38.6	51.7	25.0	Second Round	100	3.4	34.7	52	17	Rejected

(continued)

Table 2. Continued

	Results Round: 1						Results Round: 2					
	N	Mean	VC (%)	%>=4	%=5	Decision	N	Mean	VC (%)	%>=4	%=5	Decision
<ul style="list-style-type: none"> • Number of COVID-19 patients who were examined in PHC (X-ray or/and phlebotomy). Definition: Numerator: Number of COVID-19 patients who were examined in primary care (X-ray or/and phlebotomy). Denominator: Total number of patients examined in PHC (X ray or/and phlebotomy) by GP 	116	3.2	40.1	47.4	17.2	Rejected						
Indicators regarding number of procedures in PHC to COVID-19 patients												
<ul style="list-style-type: none"> • Sick leaves processed by GPs of patients in COVID-19 quarantine. Definition: Numerator: Sick leaves processed by GPs of patients in COVID-19 quarantine. Denominator: Total of sick leaves by GPs in a period 	116	3.6	35.6	58.6	33.6	Second round	100	3.6	35.4	60	32	Rejected
<ul style="list-style-type: none"> • Sick leaves processed by GPs of COVID-19 patients in isolation. Definition: Numerator: Sick leaves processed by GPs of COVID-19 patients in isolation. Denominator: Total of sick leaves by GPs in a period 	117	3.7	34.2	59.8	32.5	Second round	100	3.8	30.7	66	31	Doubtful

Low-medium scores (<70% of respondents agreed or strongly agree, and no major change was suggested for the item); medium-high scores (<70% and more than 50% in consensus ratings were sent to the second round; those with <50% in consensus ratings and without any suggestions for change by the panel after two rounds were rejected); high scores (70% of respondents agreed or strongly agreed, and at least 40% agreed to rate strongly agreed for the item).

VC: variation coefficient.

Table 3. Accepted and Uncertain Indicators based on the eDelphi Process

Indicators	Decision after the eDelphi
Indicators regarding role sentinel network	
Positive cases for SARS-CoV-2 (COVID-19) seen by the Sentinelles network.	Accepted in the first round
Positivity rates to SARS-CoV-2 (COVID-19) among all the respiratory infections by the Sentinelles network.	Accepted in the first round
Estimated incidence of COVID-19 cases per 100 000 population with respiratory signs observed in general practice through the Sentinelles network.	Accepted in the second round
Indicators regarding suspicious COVID-19 cases	
Total COVID-19 cases with positive test in primary care.	Accepted in the first round
Percentage of cases of COVID-19 among all respiratory infection cases in PHC.	Accepted in the second round
Indicators regarding primary care follow-up to COVID-19 patients	
Number of COVID-19 patients who were follow-up in primary care (nurse and/or GP) for all reasons.	Doubtful
Indicators regarding primary care follow-up to COVID-19 patients	
Number of face-to-face visits to GP with COVID-19 recorded as reason for the visit.	Accepted in the first round
Number of first visits (examinations) with COVID-19 recorded as reason for the visit (by GP)	Accepted in the second round
Indicators regarding number of procedures in PHC to COVID-19 patients	
Sick leaves processed by GPs of COVID-19 patients in isolation.	Doubtful

admissions. It is essential to establish European legislation that mandates this value, along with national regulations incorporating additional indicators to support decision-making during ARI/ILI outbreaks.

Indicators on COVID-19 cases tracing and vaccination need to be considered. When a pandemic occurs, there is a need to evaluate the impact of contact tracing to reduce virus spread. Integrating information about COVID-19 cases is crucial to establish health strategies to deal with a health emergency [2]. On the other hand, health information integrated in a same system would help to monitor vaccinations roll out and detection of possible side effect as the vaccines are still under surveillance by the European Medicines Agency (EMA). PHC is an essential place to detect potential vaccination side effects.

The high level of consensus for disaggregating data by age and vaccination status is particularly relevant in the context of a

pandemic that disproportionately affects different age groups and has seen varying rates of vaccine uptake [29]. Detailed information might help to create concrete actions to specific groups to reduce disparities and avoiding compliance of the inverse care law in PHC [30]. All the surveillance data need to be propelled back to the PHC setting and be explained how they may help in providing healthcare on a daily basis. However, during the COVID-19 pandemic many countries do not share this information [4].

Strengths and limitations

One of the main strengths lies in the participation of 31 countries, ensuring that the indicators are feasible for all participants, despite variations in the organization of PHC systems. Additionally, a high level of agreement exists for some accepted indicators, providing multiple options for monitoring the pandemic in PHC.

Limitations of this study include the potential for selection bias given the recruitment of participants with prior involvement in the Eurodata project, which may not fully represent the broader European PHC context. Additionally, the response rate, although reasonable, may not capture the full spectrum of opinions among European GPs and public healthcare professionals, particularly as the survey was only in English. The exclusion of nurses and other healthcare professionals from the panel could overlook critical insights, given their significant role in PHC, especially during the pandemic. This could be a clear limitation especially as nurses' indicators were not accepted.

While the study identifies a set of indicators, the validation of these indicators in real-world settings is not discussed. Future research should focus on testing the applicability and impact of these indicators on PHC practices and patient outcomes. Continuous evaluation and adaptation of the indicator set will be necessary to ensure they remain aligned with the current state of the pandemic and the needs of a strong PHC systems linked to public health system.

Implications for research and policy

The study's findings have the potential to significantly influence how European health authorities collect, analyze, and utilize PHC data during pandemics, improving response strategies. There is a need for subsequent studies to validate the identified indicators in various European healthcare settings. Future iterations of this study should consider including a broader range of healthcare professionals. Another option to solidify agreement could involve designing a quantitative study with a representative sample. A robust electronic reporting system is a key element. EHR systems can help collect valuable data and build indicators to measure practice workload. Health policy should be more flexible, enabling decision-making based on the epidemiological situation of ARI/ILI outbreaks. This flexibility could include hiring additional personnel, adjusting schedules, and prioritizing pathologies during periods of high workload. Unfortunately, such adaptive measures are not routinely implemented, limiting the ability of PHC to respond effectively to sudden increases in patient demand. The network of PHC providers included in surveillance reporting should be given extra support (both financial and technical) and education, be in close contact with the responsible national public health authorities and also receive feedback. Epidemiologic data should be routinely explained to PHC providers through commented reports so that the data could be used effectively to tailor health policies with regard not only to COVID-19, but ARI/ILI in general.

This eDelphi study has highlighted the complexities and variabilities inherent in the European PHC systems during the COVID-19 pandemic. The strong agreement on the need for common definitions and reproducibility of health indicators underscores the pursuit of a unified approach to pandemic data handling across Europe and the need to coordinate ECDC surveillance with PHC systems across Europe. A major challenge is the lack of national legislation for standardized PHC data collection. While some countries use EHRs and sentinel networks, many lack legal mandates, hindering disease tracking and policy decisions. A unified national PHC information system could ensure consistency and interoperability. Governments and the EU should invest in these systems and strengthen legal and technological frameworks to enhance data-driven decision-making and reinforce the role of PHC in public health preparedness and crisis response. The identified indicators can serve as a foundation for a Europe-wide PHC surveillance dashboard that could streamline data reporting, facilitate cross-country comparisons, and aid in the allocation of resources during health emergencies. Such a dashboard would also support the ECDC and WHO in their efforts to coordinate international responses to pandemics and to enhance the relationship among public health officers and PHC systems in each European country.

Conclusions

This eDelphi study contributes to a more cohesive understanding of PHC's role in pandemic surveillance and has highlighted both the potential and the challenges of standardizing PHC activity indicators for COVID-19 across Europe. The consensus reached on the selected indicators offers a pathway towards a more unified and effective surveillance system that can significantly contribute to the management of current and future pandemics, providing a foundation for the enhancement of Europe's preparedness to face them.

Authors' contributions

M.P.A.P., R.G.B., M.G.C., and S.A.B. developed the research idea. M.P.A.P. and J.J.M. carried out the body of the analysis with support of R.G.B., M.G.C., S.A.B., and T.F. M.P.A.P., R.G.B., M.G.C., S.A.B., and T.F. wrote the manuscript. All other authors contributed to the discussion and comments on the manuscript.

Acknowledgements

We would like to express our sincere gratitude to the EGPRN for their support and for providing the resources and encouragement to complete this work. We also extend our thanks to Martin Luther University Halle-Wittenberg, whose support was essential to making this work possible.

We extend our gratitude to all our colleagues for answering the questionnaires and their invaluable contributions, feedback, and unwavering support during this research project. Their participation has been instrumental in comprehensively understanding the context and deriving significant conclusions.

A.L.N. is supported by the National Institute for Health and Care Research (NIHR) Applied Research Collaboration Northwest London (NWL) and NIHR NWL Patient Safety Research Collaboration, with infrastructure support from NIHR Imperial Biomedical Research Centre. The views expressed in this publication are those of the author (s) and not necessarily those of the National Institute for Health Research or the Department of Health and Social Care.

Eurodata collaborative group: Ledia Qatipi, Elena Brutskevskaya, Steve Van den Bulck, Joris van Loenhout, Hanna Ballout, Miriam Saso, Sarah Moreels, Bohumil Seifert, Zomica Ambareva, Daniela Mileva, Veronica Rasic, Miroslav Hanzevacki, Valerija Bralic Lang, Marion Tomicic, Ivan Pristas, Zeljka Drausnik, Vasileios Trifon Karathanos, Nina Maindal, Tong Zhu, Charly Kengne Kuetche, Marion Debin, Matthieu Calafiore, Felix Bauch, Paula Tilli, Kirsi Valtonen, Christine Br€utting, Ines Lay, Sabine Klinke-Rehbein, Maria Kampouraki, Eleni Jelastopulu, Papageorgiou Dimitra Iosifina, Janos Zsuffa, Beatrix Oroszi, Peter Torzsa, Catalina Valcarcel, Lucy McShane, Hagai Levine, Alice Serafini, Austra Berzanskyte, Liga Kozlovska, Gunta Ticmane, Sabine Feldmane, Maryher Delphin Peña, Martin Sattler, Monique Aubart, Jean-Claude Leners, Marit Voltersvik, Rui Portugal, Joana Moreno, Tiago Mendes, Teresa Leão, Florentina Furtunescu, Antoaneta Dragoescu, Darinka Punosevac, Sandra Vesic Veskovac, Slavica -Dord-ovic, Natalia Enriquez Martin, Alejandra Perez Perez, Armando Chaure, Alba Gallego Royo, Ines Sebastian Sanchez, Leticia Ainhoa Sanz Asier, Francisco Rodriguez-Cabrera, Txema Coll, Naldy Parodi Lopez, Mila Gomez-Johansson, Boas Dubbelman, Fleur Otto, Jantine Woudstra, Bart Knottnerus, Ozden Gokdemir, Shushman Ivanna, Angel Gonzalez de la Fuente.

Supplementary data

Supplementary data are available at *EURPUB* online.

Conflict of interest

None declared.

Funding

This publication will be funded by open access publication fund of the Martin-Luther-University Halle-Wittenberg. This study was supported by the European General Practice Research Network (EGPRN) Grant (2022/01).

Ethical consideration

The study had the approval of the Research and Ethics Committee of the Hospital Universitario de la Paz (PI-5030), Madrid, Spain.

Data availability

The data supporting this study's findings are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

Key points

- The study identified key PHC activity indicators related to COVID-19, focusing on case detection, management, and sentinel network reporting. These indicators are intended to standardize surveillance and inform public health policy across Europe.
- Sentinel networks and electronic health records are essential for real-time data collection.
- The findings advocate for stronger support for PHC surveillance systems, including financial and technical resources, and for integrating PHC data with national public health systems to improve pandemic preparedness and management across Europe.

References

- 1 Declich S, Carter AO. Public health surveillance: historical origins, methods and evaluation. *Bull World Health Organ* 1994;72:285–304.
- 2 ECDC, WHO-Europe. Operational considerations for respiratory virus surveillance in Europe. ECDC; July 2022:1–37.
- 3 Catala M, Coma E, Alonso S *et al*. Risk diagrams based on primary care electronic medical records and linked real-time PCR data to monitor local COVID-19 outbreaks during the summer 2020: a prospective study including 7,671,862 people in Catalonia. *Front Public Health* 2021;9:693956.
- 4 Ares-Blanco S, Guisado-Clavero M, Del Rio LR *et al*. Primary care indicators for disease burden, monitoring and surveillance of COVID-19 in 31 European countries: Eurodata study. *Eur J Public Health* 2024;34:402–10.
- 5 Public Health Wales. Rapid COVID-19 Surveillance: GP Consultations for Acute Respiratory Infections. 2023 [cited 1970 Sep 16]. <https://public.tableau.com/app/profile/public.health.wales.health.protection/viz/ARIGPconsultations/Summary> (4 September 2024, date last accessed).
- 6 Ares-Blanco S, Guisado-Clavero M, Ramos Del Rio L; Clinical investigators *et al*. Clinical pathway of COVID-19 patients in primary health care in 30 European countries: Eurodata study. *Eur J Gen Pract* 2023;29:2182879.
- 7 Westerlind E, Palstam A, Sunnerhagen KS *et al*. Patterns and predictors of sick leave after Covid-19 and long Covid in a national Swedish cohort. *BMC Public Health* 2021;21:1023.
- 8 Jacob L, Koyanagi A, Smith L *et al*. Prevalence of, and factors associated with, long-term COVID-19 sick leave in working-age patients followed in general practices in Germany. *Int J Infect Dis IJID Off Publ Int Soc Infect Dis* 2021;109:203–8.
- 9 European Centre for Disease Prevention and Control (ECDC). Sentinel Surveillance. 2023 [cited 1970 Jun 28]. <https://www.ecdc.europa.eu/en/seasonal-influenza/surveillance-and-disease-data/facts-sentinel-surveillance> (4 September 2024, date last accessed).
- 10 Bagaria J, Jansen T, Marques DFP, *et al*. Rapidly adapting primary care sentinel surveillance across seven countries in Europe for COVID-19 in the first half of 2020: strengths, challenges, and lessons learned. *Euro Surveill* 2020;27:1–8.
- 11 de Lusignan S, Liyanage H, McGagh D *et al*. COVID-19 surveillance in a primary care sentinel network: in-pandemic development of an application ontology. *JMIR Public Health Surveill* 2020;6:e21434.
- 12 TESSy—The European Surveillance System Reporting Protocol for Integrated Respiratory Virus Surveillance. 2023. <https://www.ecdc.europa.eu/sites/default/files/documents/Integrated-respiratory-reporting-protocol-v1.8.pdf> (5 September 2024, date last accessed).
- 13 Hasson F, Keeney S, McKenna H. Research guidelines for the Delphi survey technique. *J Adv Nurs* 2000;32:1008–15.
- 14 McMillan SS, King M, Tully MP. How to use the nominal group and Delphi techniques. *Int J Clin Pharm* 2016;38:655–62.
- 15 Helmer-Hirschberg O. Analysis of the Future, the Delphi Method. Santa Monica, California, USA: RAND; 1967 [cited 2024 Nov 8]. (Report No.: P-3558). <http://www.rand.org/content/dam/rand/pubs/papers/2008/P3558.pdf> (4 September 2024, date last accessed).
- 16 Jünger S, Payne SA, Brine J *et al*. Guidance on Conducting and REporting DELphi studies (CREDES) in palliative care: recommendations based on a methodological systematic review. *Palliat Med* 2017;31:684–706.
- 17 European General Practice Research Network. Eurodata Collaborative group. 2021 [cited 1970 Nov 5]. <https://www.egprn.org/page/current-projects> (5 September 2024, date last accessed).
- 18 Diamond IR, Grant RC, Feldman BM *et al*. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014;67:401–9.
- 19 Guisado-Clavero M, Ares-Blanco S, Serafini A *et al*. The role of primary health care in long-term care facilities during the covid-19 pandemic in 30 European countries: a retrospective descriptive study (Eurodata study). *Prim Health Care Res Dev* 2023;24:e60.
- 20 Smith DRM, Jijón S, Oodally A *et al*. Sick leave due to COVID-19 during the first pandemic wave in France, 2020. *Occup Environ Med* 2023;80:268–72.
- 21 Coma E, Mora N, Méndez L *et al*. Primary care in the time of COVID-19: monitoring the effect of the pandemic and the lockdown measures on 34 quality of care indicators calculated for 288 primary care practices covering about 6 million people in Catalonia. *BMC Fam Pract* 2020;21:208.
- 22 Coma E, Guiriguet C, Mora N *et al*. Impact of the COVID-19 pandemic and related control measures on cancer diagnosis in Catalonia: a time-series analysis of primary care electronic health records covering about five million people. *BMJ Open* 2021;11:e047567.
- 23 Qi C, Osborne T, Bailey R *et al*. Impact of COVID-19 pandemic on incidence of long-term conditions in Wales: a population data linkage study using primary and secondary care health records. *Br J Gen Pract J R Coll Gen Pract* 2023;73:e332–9.
- 24 Moynihan R, Sanders S, Michaleff ZA *et al*. Impact of COVID-19 pandemic on utilisation of healthcare services: a systematic review. *BMJ Open* 2021;11:e045343.
- 25 Pelak M, Pettit AR, Terwiesch C *et al*. Rethinking primary care visits: how much can be eliminated, delegated or performed outside of the face-to-face visit? *J Eval Clin Pract* 2015;21:591–6.
- 26 Diercke M, Claus H, Rexroth U *et al*. Anpassung des meldesystems gemäß infektionsschutzgesetz im jahr 2020 aufgrund von COVID-19. *Bundesgesundheitsbl* 2021;64:388–94.
- 27 World Health Organization (WHO). WHO Coronavirus Disease (COVID-19) Dashboard. 2022 [cited 1970 Dec 8]. <https://covid19.who.int> (4 September 2024, date last accessed).
- 28 ECDC Europe. COVID-19: Situation Updates. 2021 [cited 1970 Dec 20]. <https://www.ecdc.europa.eu/en/covid-19> (4 September 2024, date last accessed).
- 29 WHO. Covid-19 Strategic Preparedness; February 2021. https://www.who.int/docs/default-source/coronaviruse/covid-strategy-update-14april2020.pdf?sfvrsn=29da3ba0_19 (2 September 2024, date last accessed).
- 30 Hart JT. The inverse care law. *Lancet London Engl* 1971;1:405–12.

© The Author(s) 2025. Published by Oxford University Press on behalf of the European Public Health Association.
This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com
European Journal of Public Health, 2025, 35, 1026–1035
<https://doi.org/10.1093/eurpub/ckaf051>
Original Manuscript