

This is the peer reviewed version of the following article:

Functional Outcome of Elderly Patients Treated for Odontoid Fracture: A Multicenter Study / De Bonis, P.; Iaccarino, C.; Musio, A.; Martucci, A.; De Iure, F.; Donati, R.; Cultrera, F.; Tosatto, L.; Servadei, F.; Alesi, D.; Cavallo, M. A.; Ghadirpour, R.; Molinari, F.; Lofrese, G.. - In: SPINE. - ISSN 0362-2436. - 44:13(2019), pp. 951-958. [10.1097/BRS.0000000000002982]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

04/05/2026 03:03

(Article begins on next page)

Spine (Phila Pa 1976)

Functional outcome of elderly patients treated for odontoid fracture. A multicenter study. --Manuscript Draft--

Manuscript Number:	SPINE 157299R2
Full Title:	Functional outcome of elderly patients treated for odontoid fracture. A multicenter study.
Article Type:	Cervical Spine
Keywords:	C2, odontoid, fracture, comorbidity, disability, elderly, ultra-elderly, functional outcome, stability, healing, treatment, fusion, surgery, conservative, crossover
Corresponding Author:	Giorgio Lofrese, M.D. Bologna, Italy ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	
Corresponding Author's Secondary Institution:	
First Author:	Pasquale De Bonis, Prof, Ph.D
First Author Secondary Information:	
Order of Authors:	Pasquale De Bonis, Prof, Ph.D
	Corrado Iaccarino, M.D.
	Antonio Musio, M.D.
	Antonio Martucci, M.D.
	Federico De Iure, M.D.
	Roberto Donati, M.D.
	Francesco Cultrera, M.D.
	Luigino Tosatto, M.D.
	Franco Servadei, Prof
	Domenico Alesi, M.D.
	Michele Alessandro Cavallo, Prof
	Reza Ghadirpour, M.D.
	Filippo Molinari, M.D.
Giorgio Lofrese, M.D.	
Order of Authors Secondary Information:	
Additional Information:	
Question	Response
Please provide the Word Count of your manuscript text. Include only the main body of text (exclude abstract, references, figures, and table legends).	2698
Please provide the Word Count of your structured abstract. Include only the	295

abstract, not key words.	
Please select the level of evidence for this manuscript.	3
<p>Device Status/Drug Statement: Please select the statement below that applies to your submission. After you have selected the appropriate statement, <u>Please also add the same statement to the Title Page of your submission.</u></p>	The legal regulatory status of the device(s)/drug(s) that is/are the subject of this manuscript is not applicable in my country.
<p>RETAINED RIGHTS: Except for copyright, other proprietary rights related to the Work (e.g., patent or other rights to any process or procedure) shall be retained by the author. To reproduce any text, figures, tables, or illustrations from this Work in future works of their own, the author must obtain written permission from Wolters Kluwer Health, Inc. ("WKH").</p> <p>ORIGINALITY: Each author warrants that his or her submission to the Work is original, does not infringe upon, violate, or misappropriate any copyright or other intellectual property rights, or any other proprietary right, contract or other right or interest of any third party, and that he or she has full power to enter into this agreement. Neither this Work nor a similar work has been published nor shall be submitted for publication elsewhere while under consideration by this Publication.</p> <p>AUTHORSHIP RESPONSIBILITY: Each author warrants that he or she has participated sufficiently in the intellectual content, the analysis of data, if applicable, and the writing of the Work to take public responsibility for it. Each has reviewed the final version of the Work, believes it represents valid work, and approves it for publication. Moreover, should the editors of the Publication request the data upon which the work is based, they shall</p>	I agree

produce it.

PREPRINTS: Upon acceptance of the article for publication, each author warrants that he/she will promptly remove any prior versions of this Work (normally a preprint) that may have been posted to an electronic server.

DISCLAIMER: Each author warrants that this Work contains no libelous or unlawful statements and does not infringe or violate the publicity or privacy rights of any third party, libel or slander any third party, contain any scandalous, obscene, or negligently prepared information, or infringe or violate any other personal or proprietary right of others. Each author warrants that the Work does not contain any fraudulent, plagiarized or incorrectly attributed material. Each author warrants that all statements contained in the Work purporting to be facts are true, and any formula or instruction contained in the Work will not, if followed accurately, cause any injury, illness, or damage to the user. If excerpts (e.g., text, figures, tables, illustrations, or audio/video files) from copyrighted works are included, a written release will be secured by the author prior to submission, and credit to the original publication will be properly acknowledged. Each author further warrants that he or she has obtained, prior to submission, written releases from patients whose names or likenesses are submitted as part of the Work. Should the Editor or WKH request copies of such written releases, the author shall provide them in a timely manner.

DISCLOSURES/CONFLICT OF INTEREST

Each author must identify any financial interests or affiliations with institutions, organizations, or companies relevant to

<p>the manuscript by completing the form below. Additionally, any financial associations involving a spouse, partner or children must be disclosed as well.</p> <p>Note: Some sections below come from the ICMJE Uniform Disclosure Form for Potential Conflicts of Interest at http://www.icmje.org/downloads/coi_disclosure.pdf (dated July 2010).</p>	
<p>Did you or your institution at any time receive payment or support in kind for any aspect of the submitted work (including but not limited to grants, consulting fee or honorarium, support for travel to meetings for the study or other purposes, fees for participation in review activities such as data monitoring boards, statistical analysis, end point committees, and the like, payment for writing or reviewing the manuscript, provision of writing assistance, medicines, equipment, or administrative support, etc...)?</p>	<p>No</p>
<p>Other: Did you or your institution at any time receive additional payments or support in kind for any aspect of the submitted work?</p>	
<p>Please indicate whether you have financial relationships (regardless of amount of compensation) with entities. You should report relationships that were present during the 36 months prior to submission including board membership, consultancy, employment, expert testimony, grants/grants pending, payment for lectures including service on speakers bureaus, payment for manuscript preparation, patents (planned, pending or issued), royalties, payment for development of educational presentations, stock/stock options, travel/accommodations/meeting expenses unrelated to activities listed (for example, if you report a consultancy above there is no need to report travel related to that consultancy), etc.</p>	<p>No</p>
<p>Other (err on the side of full disclosure):</p>	<p>No additional financial relationships</p>

<p>Please indicate whether you have any additional financial relationships (regardless of amount of compensation) with entities. You should report relationships that were present during the 36 months prior to submission.</p>	
<p>Other Relationships</p> <p>Are there other relationships or activities that readers could perceive to have influenced, or that give the appearance of potentially influencing, what you wrote in the submitted work?</p>	<p>No other relationships/conditions/circumstances that present potential conflict of interest</p>
<p>AUTHOR'S OWN WORK: In consideration of WKH's publication of the Work, the author hereby transfers, assigns, and otherwise conveys all his/her copyright ownership worldwide, in all languages, and in all forms of media now or hereafter known, including electronic media such as CD-ROM, Internet, and Intranet, to WKH. If WKH should decide for any reason not to publish the Work, WKH shall give prompt notice of its decision to the corresponding author, this agreement shall terminate, and neither the author nor WKH shall be under any further liability or obligation. Each author grants WKH the rights to use his or her name and biographical data (including professional affiliation) in the Work and in its or the journal's promotion. Notwithstanding the foregoing, this paragraph shall not apply, and any transfer made pursuant to this paragraph shall be null and void if (i) the Work has been accepted by WKH for publication, and (ii) the author chooses to have the Work published by WKH as an open access publication.</p> <p>WORK MADE FOR HIRE: If this Work or any element thereof has been commissioned by another person or organization, or if it has been written as part of the duties of an employee, an authorized representative of the commissioning organization or employer must also sign this form stating his or her title in the organization.</p>	<p>I agree</p>

GOVERNMENT EMPLOYEES: If the Work or a portion of it has been created in the course of any author's employment by the United States Government, check the "Government" box at the end of this form. A work prepared by a government employee as part of his or her official duties is called a "work of the U.S. Government" and is not subject to copyright. If it is not prepared as part of the employee's official duties, it may be subject to copyright.

INSTITUTIONAL REVIEW BOARD/ANIMAL CARE COMMITTEE APPROVAL: Each author warrants that his or her institution has approved the protocol for any investigation involving humans or animals and that all experimentation was conducted in conformity with ethical and humane principles of research.

WARRANTIES: Each author warranty made in this form is for the benefit of WKH and the Editor; each author agrees to defend, indemnify, and hold harmless those parties for any breach of such warranties.

Spine will permit the author(s) to deposit for display a "final peer-reviewed manuscript" (the final manuscript after peer-review and acceptance for publication but prior to the publisher's copyediting, design, formatting, and other services) 12 months after publication of the final article on the author's personal web site, university's institutional repository or employer's intranet, subject to the following:

* You may only deposit the final peer-reviewed manuscript.

* You may not update the final peer-reviewed manuscript text or replace it with

I agree

<p>a proof or with the final published version.</p> <p>* You may not include the final peer-reviewed manuscript or any other version of the article on any commercial site or in any repository owned or operated by any third party. For authors of articles based on research funded by the National Institutes of Health ("NIH"), Wellcome Trust, Howard Hughes Medical Institute ("HHMI"), or other funding agency, see below for the services that WKH will provide on your behalf to comply with "Public Access Policy" guidelines.</p> <p>* You may not display the final peer-reviewed manuscript until twelve months after publication of the final article.</p> <p>* You must attach the following notice to the final peer-reviewed manuscript: "This is a non-final version of an article published in final form in (provide complete journal citation)".</p> <p>* You shall provide a link in the final peer-reviewed manuscript to the Spine website.</p>	
<p>"Public Access Policy" Funding Disclosure Please disclose below if you have received funding for research on which your article is based from any of the following organizations:</p>	
<p>Please select:</p>	<p>Author's Own Work</p>
<p>Any additional comments?</p>	
<p><u>Compliance with RCUK and Wellcome Trust Open Access Policies</u></p> <p>Both the Research Councils UK (RCUK) and the Wellcome Trust have adopted policies regarding Open Access to articles that have been funded by grants from the RCUK or the Wellcome Trust. If either "Wellcome</p>	<p>I agree</p>

Trust” or “Research Councils UK (RCUK)” has been selected above, and the authors of the applicable article choose to have the article published as an open access publication, the following policies will apply:

* If the article is to be published pursuant to the “Gold” route of Open Access, both the RCUK and the Wellcome Trust require that WKH make the article freely available immediately pursuant to the Attribution 4.0 Creative Commons License, currently found at <http://creativecommons.org/licenses/by/4.0/legalcode> (the “CC BY License”). The CC BY License is the most accommodating of the Creative Commons licenses and allows others to distribute, remix, tweak, and build upon the article, even commercially, as long as they credit the authors for the original creation.

* If the article is to be published pursuant to the “Green” route of Open Access, both the RCUK and the Wellcome Trust require that WKH make the article freely available within six months pursuant to the Attribution-NonCommercial 4.0 Creative Commons License, currently found at <http://creativecommons.org/licenses/by-nc/4.0/legalcode> (the “CC BY-NC License”). The CC BY-NC License allows others to remix, tweak, and build upon the article non-commercially, and although their new works must also acknowledge the authors for the original creation and be non-commercial, they don't have to license their derivative works on the same terms.

As a service to our authors, WKH will identify the National Library of Medicine (NLM) articles that require deposit pursuant to the RCUK and Wellcome

<p>Trust policies described in this section. This Copyright Transfer Agreement provides the mechanism for identifying such articles.</p> <p>WKH will transmit the final peer-reviewed manuscript of an article based on research funded in whole or in part by either RCUK or the Wellcome Trust to Pub Med Central.</p> <p>Upon NIH request, it remains the legal responsibility of the author to confirm with NIH the provenance of his/her manuscript for purposes of deposit. Author will not deposit articles him/herself. Author will not alter the final peer-reviewed manuscript already transmitted to NIH.</p> <p>With respect to the “Green” route of Open Access, author will not authorize the display of the final peer-reviewed manuscript prior to 6 months following publication of the final article.</p> <p>Authors of articles that have been funded from grants from the RCUK or the Wellcome Trust are required to sign the WKH Open Access License Agreement prior to publication of the applicable article. Please contact the Editorial Office of the applicable journal to receive the Open Access License Agreement that is to be signed in connection with the publication of the article.</p>	
<p>I am the person in question for this submission or otherwise have approval to complete this agreement.</p>	<p>I agree</p>
<p>CME/CE Disclosure</p> <p>Each author must identify and disclose any financial associations involving a spouse, partner or children by completing</p>	<p>I agree</p>

<p>the Family Disclosure question below, and whether any off-label uses or unapproved drugs or devices are discussed in his/her manuscript by completing the Off-Label Use/Unapproved Drugs or Products question below. In the event that the Work is published as a continuing education or continuing medical education article, this information will be provided to the accrediting body and may be included in the published article. When applicable, articles accepted for publication may need to comply with additional standards related to CME or CE accreditation. Please refer to guidelines for authors for details.</p> <p>WKH and its affiliates reserve the right to publish the manuscript as a continuing education article.</p>	
<p>Family Disclosure</p> <p>Do your children or your spouse or partner have financial relationships with entities that have an interest in the content of the submitted work?</p>	<p>No other relationships/conditions/circumstances that present potential conflict of interest</p>
<p>Off-Label Use/Unapproved Drugs or Products</p> <p>If your manuscript discusses an unlabeled use of a commercial product or device or an investigational use of a product or device not yet approved by the FDA for any purpose, you must specifically disclose in the manuscript that the product is not labeled for the use under discussion or that the product is still investigational. Please check the item below that applies to you</p>	<p>I will not discuss unlabeled/investigational uses of any commercial product or device</p>

Reviewers' comments:

Reviewer #1: Abstract:

More data has to be included: Please add how many patients have been treated by surgery (and by which technique) compared to conservative treatment.

We have integrated the abstract according to your suggestion.

Materials and methods:

"Type III odontoid fractures were considered unstable whenever a shallow-type, an antero-posterior displacement of more than 5mm and an odontoid angulation of more than 11° was noted."

Did you consider all type 2 fractures as unstable?

Thank you for your question. All type II fractures have been considered potentially unstable and, since those ones revealing an antero-posterior displacement over 5mm should be considered severely unstable, we have included this value in the multivariate analysis, in order to establish if "severely unstable" fractures may have a different outcome. Therefore, all the patients included in the study had an unstable odontoid fracture, because of the radiological evidence of a type II or alternatively of a type III shallow-type/A-P dislocated >5mm/angulated >11°. Manuscript has been modified accordingly

Treatment modalities:

Basically, patients with stable fractures have been treated non-operatively (cervical collar or halo vest), and all other patients have been treated by one of three different operative techniques. If this is how patients have been treated you should emphasize this (in abstract and text). I would strongly recommend that you provide data with exact numbers of patients treated by halo vs cervical collar for a stable fracture (how many non-fusions, how many cross-overs, functional outcome) and of patients treated by the three operative techniques for unstable fractures and compare the outcome of these techniques.

Thank you for your comment. Since all type II odontoid fractures have been considered as unstable, the choice of treatment (conservative Vs surgical) appeared influenced only by the antero-posterior displacement (< or >5mm) or by the surgeon decision: in fact, in some cases, despite a higher antero-posterior displacement, the surgeon decided to treat patients with collar or halo, due to patient clinical conditions that would have increased the risk of surgery unacceptably. In order to correct this selection bias, we performed a multivariate analysis including Comorbidity indexes (Charlson, ASA, Modified Rankin Scale). Other radiological fracture characteristics had no role in setting treatment strategy. As suggested, we've modified abstract and manuscript in order to emphasize this concept. According to your recommendation, but respecting the aforementioned assumption of instability for type II fractures, we've integrated the manuscript with the most significant data regarding functional outcome and cross-over for all the treatments adopted (conservative and surgical), comparing them in the discussion. Since all the 25 "cross-over" patients needed for a second step treatment because of a mobile non-fusion at the dynamic cervical spine imaging, non-fusion data should be discussed including this subgroup of patients, which has been excluded from the analysis from the beginning in order to avoid confusion. According to the previous revision requested, cross-over has been considered as an outcome parameter, preserving the main purpose of our study which is the analysis of functional outcome. Functional outcome for patients with crossover cannot be evaluated properly because it is impossible to establish if outcome is more related to initial treatment or second treatment or both. For this reason, like in all studies analyzing outcome, the subgroup of patients with treatment crossover was not included in the analysis of functional outcome. These patients would deserve a detailed method, another stat, results, tables and discussion rather than a paragraph in this manuscript, which would risk to generate confusion in the reader considering the topic of this study.

The advantage of your investigation is that you obtained clinical outcome parameters for a large number of patients.

The results of your investigation show what is already known: Stable fractures do have a better outcome than unstable fractures. Anterior screw fixation in elderly patients shows higher rates of re-operation.

Thank you for your comment. As you noted cross-over regarded even patients who underwent surgery and more specifically 3/25 patients needed for a second step C1-C2 posterior arthrodesis as a rescue procedure after mechanical failure of anterior odontoid screw fixation. The manuscript has been integrated with this data.

Functional outcome of elderly patients treated for odontoid fracture. A multicenter study.

Pasquale De Bonis*¹, Prof., Ph.D., M.D., Corrado Iaccarino*², M.D., Antonio Musio*¹, M.D., Antonio Martucci³, M.D., Federico De Iure³, M.D., Roberto Donati⁴, M.D., Francesco Cultrera⁴, M.D., Luigino Tosatto⁴, M.D., Franco Servadei², Prof., M.D., Domenico Alesi³, M.D., Michele Alessandro Cavallo¹, Prof., M.D., Reza Ghadirpour², M.D., Filippo Molinari², M.D., Giorgio Lofrese*⁴, M.D.

¹Neurosurgery Division, University Hospital S.Anna, Cona di Ferrara, Italy

²Neurosurgery-Neurotraumatology Division, University Hospital of Parma, Parma, Italy

³Department of Spine Surgery, Maggiore “C.A. Pizzardi” Hospital, Bologna, Italy

⁴Neurosurgery Division, “M. Bufalini” Hospital, Cesena, Italy

Corresponding Author:

Giorgio Lofrese*², M.D.

Neurosurgery Division, “M. Bufalini” Hospital

Viale Ghirotti 286, 47521 Cesena - Italy

e-mail: giorgio.lofrese@gmail.com

tel: +39 328 05 82 502

fax: +39 0547 352918

on behalf of the “DENSER group”

*these authors equally contributed to the paper

The legal regulatory status of the device(s)/drug(s) that is/are the subject of this manuscript is not applicable in my country.

No funds were received in support of this work.

Relevant financial activities outside the submitted work: consultancy, grants.

ABSTRACT

Study Design: Retrospective multicenter study.

Objective: Analysis of impact of conservative and surgical treatments on functional outcome of geriatric odontoid fractures.

Summary of Background Data: Treatment of odontoid fractures in aged population is still debatable.

Methods: 147 consecutive odontoid fractures in elderly patients were classified according to Anderson-D'Alonzo and Roy-Camille classifications. Philadelphia type collar was always positioned and kept as a treatment whenever acceptable. Halo-vest, anterior screw fixation, C1-C2 posterior arthrodesis and occipito-cervical fixation were the other treatments adopted. Conservative or surgical treatment strategy were more significantly influenced by antero-posterior displacement ($<$ or $>$ 5 mm) and by surgeon decision. On admission ASA, modified Rankin scale (mRS-pre) and Charlson Comorbidity Index (CCI) were assessed. Modified Rankin scale (mRS-post), Neck Disability Index (NDI) and Smiley Webster Pain Scale (SWPS) were administered 12-15 months after treatment to estimate functional outcome in terms of general disability, neck-related disability and ability to return to work/former activity. Risk of treatment crossover was calculated considering factors affecting outcome. Fracture healing process in terms of fusion-stability, no fusion-stability, no fusion-no stability was evaluated at 12 months through a cervical CT scan. Dynamic cervical spine x-rays were obtained whether necessary. No fusion-stability was considered an adequate treatment goal in our geriatric population. Chi square/Fisher exact test and logistic regression were performed for statistical analysis.

Results: Overall 67 patients were treated conservatively whereas 80 underwent surgery. Collar was adopted in 45 patients, while anterior odontoid fixation and C1-C2 posterior arthrodesis were preferred for 30 patients each. 79.8% of patients showed good outcomes according to NDI. No significant differences were observed between patients of 65-79 years and \geq 80 years ($p=0.81$). CCI greatly correlated with mRS-post, with higher indexes in 68.8% of cases characterized by good outcomes ($p=0.05$). mRS-pre correlated with NDI ($p<0.000001$) and mRS-post ($p=0.04$). CCI, mRS-pre and surgery were associated with worse NDI, while both C1-C2 posterior arthrodesis and occipito-cervical stabilization were associated with worse mRS-post, respectively in 40% and 30% of cases. Younger patients had a higher risk of treatment crossover.

Conclusions: mRS-pre and CCI provided two independent predictive values respectively for functional outcome and post-treatment disability. Compared to conservative immobilizations, surgery revealed no advantages in the elderly in terms of functional outcome.

Keywords: C2, odontoid, fracture, comorbidity, disability, elderly, ultra-elderly, functional outcome, stability, healing, treatment, fusion, surgery, conservative, crossover

Abbreviations: mRS-pre = pre-treatment modified Rankin scale; mRS-post = post-treatment modified Rankin scale; CCI = Charlson Comorbidity Index; NDI = Neck Disability Index; SWPS = Smiley Webster Pain Scale; MPR = multiplanar reconstruction;

Level of Evidence: 3

Keypoints

- Outcome of odontoid fractures should not be measured with fusion rate, but with functional outcome scoring systems;
- Pre-treatment higher comorbidity rate (Charlson Comorbidity Index) and higher disability (Modified Rankin Scale) are associated with worse functional outcome in geriatric type II odontoid fractures regardless of treatment;
- Surgery is related to higher risk of worse functional outcome in patients aged 65 or over compared with external immobilization with collar or halo.

Mini Abstract

Functional outcome of elderly patients treated for odontoid fracture is strictly related to their pre-injury level of function and comorbidity. Compared to conservative external immobilizations, surgery revealed no advantages in the elderly in terms of functional outcome.

INTRODUCTION

1
2
3
4
5 Odontoid fractures represent the most common cervical spine fractures among elderly and the
6
7 most frequent vertebral fractures among patients over 80 years. Their rate in this old patient
8
9 group is increasing, growing faster than the ageing population¹.

10
11
12 Despite recent advances in internal fixation techniques, controversy remains regarding whether
13
14 operative or non-operative management is the best treatment approach for elderly patients²⁻¹¹.

15
16 Non-operative treatment options include hard cervical collar and halo-vest orthosis, while
17
18 operative strategies comprehend anterior odontoid osteosynthesis, C1-C2 posterior arthrodesis
19
20 and occipito-cervical stabilization. Treatments with cervical immobilization avoid the risks of
21
22 surgery but they are associated with higher non-union rates, whereas surgical fixations put
23
24 patients at risk for post-operative complications but leading to significantly better union rates¹.

25
26
27 The outcomes for those patients showing nor fusion nor signs of fracture healing process remain
28
29 unclear and morbidity of odontoid pseudoarthrosis in aged population is also not well
30
31 defined^{12,13}. Many factors have to be taken into account to find the right balance between
32
33 support of fracture consolidation and risk of treatment complications. Based on these
34
35 considerations, the decision for either conservative or surgical therapy is made^{11,14}. In this
36
37 particular subset of patients, most studies focus on radiological outcome and survival, without
38
39 considering how different treatment modalities may have an impact on daily life^{3,5,12,14-19}. The
40
41 aim of this study was to assess the functional outcome of elderly patients with unstable odontoid
42
43 fractures in relation to the different treatment modalities, focusing on the impact of external
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

immobilizations, surgical fixations and comorbidities in such a fragile population.

MATERIALS AND METHODS

Study population and radiological classification

From January 2012 through December 2016, 210 consecutive patients treated for unstable odontoid fractures in four different trauma centres were included in this retrospective analysis.

Three different causes of injury were identified: vehicle accidents (including any kind of means of transport, or as a pedestrian); falls with low energy impact (less than 2 meters); falls with high energy impact (more than 2 meters). Nine patients died, two were lost at follow-up and those ones under 65 years or with missing/incomplete data were excluded from the study. Six patients with spinal cord injury were also excluded because their functional outcome would have been affected by their neurological impairment. Among the remaining 172 patients, 25 were excluded because of treatment crossover. All the 147 enrolled patients underwent a baseline cervical spine CT evaluation at the hospital admission and at 1, 6 and 12 months after surgery as part of a standardized radiological follow-up.

CT scans of the cervical spine were performed using a clinical multidetector scanner. Images always contained multiplanar reconstructions (MPR), which were systematically reviewed to classify each fracture according to the Anderson and D'Alonzo (type II or III) and the Roy-Camille classifications (anterior oblique, posterior oblique and horizontal) respectively describing fracture localization and direction of fracture line. All type II odontoid fractures have been considered potentially unstable and those ones, among them, revealing an antero-posterior displacement over 5 mm, have been considered potentially severely unstable. Type III odontoid

fractures were considered unstable whenever a shallow-type, an antero-posterior displacement of more than 5 mm and an odontoid angulation of more than 11° was noted.

Treatment modalities

All the patients had a Philadelphia® type cervical collar (Center for Prosthetics Orthotics, Inc., Seattle, WA, USA) positioned soon after the spine trauma and worn as an acceptable treatment or as a temporary immobilization before surgery, depending on the decisions taken by the physicians after the imaging of the cervical spine. No protocols were applied to drive the choice of the most appropriate strategy and patients were managed conservatively or surgically according to knowledge and experience of dedicated neurosurgeons/orthopedics and their surgical armamentarium. However, all the treating physicians preferred surgery whenever a fracture gap of more than 2 mm, an antero-posterior displacement of more than 5 mm and an odontoid angulation of more than 11° occurred. Conservative or surgical treatment strategy were more significantly influenced by antero-posterior displacement (< or >5 mm) and by surgeon decision. More specifically, despite an antero-posterior displacement over 5 mm, in some cases surgeon decided to treat patients with external immobilization, due to patient clinical conditions that would have unacceptably increased the risk of surgery. Other radiological fracture characteristics played no role in setting treatment strategy. Five different treatments were reported: hard cervical collar (Philadelphia® [Center for Prosthetics Orthotics, Inc., Seattle, WA, USA], Aspen® or Vista® [Aspen Medical Products, Irvine, CA, USA] type) and halo vest (PMT® Corporation, Chanhassen, MN, USA) among those ones conservatives; anterior odontoid screw fixation, C1-C2 posterior arthrodesis (Harms technique) and occipito-cervical stabilization among the surgical options. Risk of treatment crossover was calculated for all the factors affecting outcome, but those patients who needed for such a second step surgery were excluded from the analysis due to the impossibility to establish the role of first,

second or both treatments in determining the final functional outcome.

Functional and radiological assessment

On admission, each patient was clinically assessed adopting ASA score, modified Rankin scale (mRS-pre) and Charlson Comorbidity Index (CCI) respectively for estimating general physical status, degree of disability and mortality risk according to comorbidities (*table 1*).

From 12 to 15 months after treatment, functional evaluations were performed employing a second modified Rankin scale (mRS-post) together with the Neck Disability Index (NDI) and the Smiley Webster Pain Scale (SWPS), investigating respectively general disability, neck-related disability and ability to return to work/former activity. Both NDI and SWPS were delivered as phone interview questionnaires by two different operators whether necessary.

The radiological outcome was evaluated through CT scans of the cervical spine with MPR 12 months after treatment. In cases with doubtful fracture healing attitude, dynamic cervical spine x-rays were obtained to rule out atlantoaxial instability. According to the evidences of both CT scan and dynamic x-rays, three different conditions were identified: fusion-stability, no fusion-stability, no fusion-no stability.

Fusion was defined by the evidence of bone trabeculae crossing the fracture line in absence of sclerotic borders/bone resorption of fracture's fragments. Fracture stability was determined by the absence of secondary displacement of the odontoid process. This latter condition was considered an adequate treatment goal in our geriatric population, even without radiological evidences of bony fusion (no fusion-stability).

All the surgical consents included paragraphs regarding patient agreement to elaborate and share disease-related information for scientific purposes. A similar agreement form, included

1 in a paper with indications for management of external immobilization, was signed at first
2 follow-up appointment by those patients non-operatively treated.
3
4
5

6 7 **Statistical analysis**

8
9 The statistical analyses were performed with the Statistical Package for the Social Sciences
10 software for Windows (SPSS Inc. Released 2008. SPSS Statistics for Windows, Version 17.0.
11 Chicago, IL, USA). Univariate analysis included impact of age (65-79 vs 80 and over),
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
The statistical analyses were performed with the Statistical Package for the Social Sciences software for Windows (SPSS Inc. Released 2008. SPSS Statistics for Windows, Version 17.0. Chicago, IL, USA). Univariate analysis included impact of age (65-79 vs 80 and over), Charlson Comorbidity Index (2-5 vs 6-8), ASA score (1-2 vs 3 vs 4), mRS-pre (0-3 vs 4-5), Anderson-D'Alonzo classification (odontoid fracture type II vs III vs II-III plus other cervical fractures), fragment dislocation (more or less than 5 mm) and treatment (collar vs halo vs anterior odontoid screw vs C1-C2 posterior arthrodesis vs occipito-cervical stabilization) on outcome.

Outcome variables were analyzed: complications (yes vs no), radiological outcome (fusion, no fusion-stable, no fusion-unstable), NDI (0-28% vs 30-48% vs 50-100%), mRS-post (0-3 vs 4-5), SWPS (excellent or good vs fair or poor) and crossover (yes vs no).

Chi square/Fisher exact test were adopted to compare variables. Logistic regression analysis investigated the impact of the aforementioned variables on dichotomous outcome variables, including NDI 1-48% vs 50-100%. Results presenting $p < 0.05$ were considered statistically significant.

68 69 **Source of funding**

70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
The authors report no conflict of interest concerning the materials or methods used in this study or the findings specified in this paper. No funds or grants have been received for this study.

RESULTS

Among all the 147 patients (73 male, 74 female) included in this multicenter study, 51.7% were aged 80 or over. In 71.4% of the cases the odontoid fracture derived from a fall with low energy impact, while in 19% occurred for a vehicle accident. Type II fractures were recorded in 58.6% of the cases, while in 27.9% type II or type III fractures were associated with other lesions of the craniocervical junction. Only 8% of the patients had an odontoid process dislocation ≥ 5 mm.

Treatment

Most patients treated conservatively (45.6% of total), received a Philadelphia type hard cervical collar, while patients who underwent surgery (54.4% of total) were treated as follows: anterior odontoid screw fixation (20.4%), C1-C2 posterior arthrodesis (20.4%), and occipito-cervical stabilization (13.6%). Overall treatment-related complications were 5.3% over 80 years and 7% between 65 and 79 years (Pearson's Chi Square 2-sided, $p=0.65$). Correlating the different treatment modalities with NDI, a good outcome (NDI 0-48%) was recorded in: 100% of hard cervical collar, 95.5% of halo vest, 90% of occipito-cervical stabilization, 47.6% of anterior odontoid screw fixations and 45.5% of C1-C2 posterior arthrodesis (Pearson chi squared 2 sided, $p<0.0001$). Similarly, a positive outcome, in terms of SWPS excellent or good, was reported in 88.9%, 81.8% and 75% of patients respectively treated with collar, halo vest and occipito-cervical stabilization, and only in 47.6% and 59.1% of those who respectively underwent anterior odontoid screw fixation and C1-C2 posterior arthrodesis (Pearson's Chi

Square 2-sided, $p < 0.0001$).

Radiological outcome

At 12-month follow up, 54% of patients obtained a complete fracture healing process while 36.5% didn't reach the fusion although revealing stable at dynamic x-rays. No significant differences were observed in the rate of fusion between patients over 80 and those ones between 65 and 79 years (Pearson's Chi Square 2-sided, $p = 0.74$)

Functional outcome

According with NDI, 79.8% of patients had a good outcome (range 0-48%). No differences were observed between younger (65-79 years) and older (≥ 80 years) patients (Pearson's Chi Square 2-sided, $p = 0.81$).

Accordingly, the outcome based on SWPS showed a 48.5% of good and 23.8% of excellent outcome in terms of return to work/former activities with a post-treatment level of satisfaction equally outstanding in both age subgroups (Fisher's exact test 2-sided, $p = 0.56$).

A CCI from 2 to 5 and an ASA score of 3 were recorded respectively in 66% and 63.9% of patients, and most of them needed no assistance neither before or after treatment with mRS-pre and mRS-post ranging from 0 to 3 respectively in 70.1% and 76.2% of cases.

As expected, CCI significantly correlated with mRS-post, with higher indexes (6-8) reported in 68.8% of cases characterized by worse outcomes (Fisher's exact test 2-sided, $p = 0.05$).

Pre-treatment mRS strongly correlated with NDI (92% of patients with pre-treatment mRS 0-3 had a NDI between 0% and 48%-Fisher's exact test. 2-sided, $p < 0.0001$) and with post-treatment mRS (83.3% of patients with pre-treatment mRS 0-3 had a post-treatment mRS 0-3, Fisher's exact test 2-sided, $p = 0.04$).

Patient's satisfaction was reported as excellent or good (SWPS) in 83.2% of patients with mRS-

pre 0-3, but only in 41.4% of patients with mRS-pre 4-5 (Fisher's Exact test 2-sided, $p < 0.0001$).

No correlation between mRS-pre and radiological outcome was observed.

Factors affecting functional outcome

When correcting for confounding variables at logistic regression analysis, we found that high values of CCI and mRS-pre, together with all surgical procedures (anterior odontoid screw fixation, C1-C2 posterior arthrodesis and occipito-cervical stabilization) were associated with worse NDI (*Figure 1, table 2*). Likewise, high scores of mRS-pre and surgery were associated with worse SWPS (*table 3*). Multivariate analysis also showed that C1-C2 posterior arthrodesis and occipito-cervical stabilization were associated with worse mRS-post, respectively in 40% and 30 % of cases (*table 4*), while C1-C2 posterior arthrodesis and ultra-elderly patients were associated with a lower risk of crossover (*table 5*). Among the 25 patients who needed a treatment crossover, thus not included in our outcome analysis, 13 were wearing a halo vest and 9 a Philadelphia type collar. The remaining 3 patients underwent C1-C2 posterior arthrodesis as rescue surgery for loosening of the anterior odontoid screw.

DISCUSSION

To date, this multicenter study represents one of the largest investigations on odontoid fractures and the largest study on functional outcome in the elderly.

The main limitations of this study lie in its retrospective nature, in the number of dead patients and in the presence of patients needing a treatment crossover. Nonetheless, the percentage of patients with missing data or lost at follow-up (15 out of 204-7.3%) and the percentage of patients with treatment crossover (25 out of 172-14.5%) is relatively low.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Discordance between radiological and functional outcome can be explained with the particular fragility of this subset of patients expressed by the poor post-surgery level of function: sharp dissection of paraspinal muscles and ligaments together with the fixation itself have both an influence on outcome much more than fracture union. On the other hand, such geriatric population shows an age-related cervical spondylosis, which represents a pre-existent limitation likely not overloading any eventual odontoid fibrous union, because of the reduced range of motion of head and neck usually associated to this degenerative spine condition.

Interestingly, according to the scales used for the assessment of functional outcome, patients aged 80 years or over had no worse results than those between 65 and 79 years-old, with these latter revealing even a higher risk of crossing over from conservative to surgical treatment.

Independently from the age group, multivariate analysis showed that greater numbers of comorbidities, thus higher CCIs, were related to more severe degrees of disability.

The mRS-pre represented a strong independent predictive value for functional outcome.

As expected, independently from the conservative/surgical strategy, patients needing assistance before treatment (i.e. with higher mRS-pre values) didn't improve their post-treatment score, therefore indicating a marginal role of surgery whenever a severe disability already exists. On this line, patients with mRS-pre 4 or 5 revealed a favourable NDI (0-48%) and an excellent or good SWPS only in 37.9% and 41.4% of cases, respectively. Patients with a low mRS-pre (0-3), instead, had a favourable NDI (0-48%) and an excellent or good SWPS in 92% and 83.2% of cases, respectively.

Such deep difference expressed in percentages demonstrated both the effectiveness and reliability of mRS-pre in estimating the functional outcome, confirming what assumed by Vaccaro et al. ¹¹ about the pre-injury level of function and its relevant impact on outcome regardless the treatment type.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Logistic regression analysis showed that CCI, mRS-pre and all the surgical procedures were associated with worse NDI while both C1-C2 posterior arthrodesis and occipito-cervical stabilization were associated with worse mRS-post. Differently from Joestl et al. ⁶, who observed better functional results among patients who underwent anterior odontoid screw fixation compared with those ones treated with halo, we noticed a homogeneously worse functional outcome after any kind of surgical procedure, but an expectable lower risk of crossover after C1-C2 posterior arthrodesis. The risk (OR) of having a worse NDI after surgery was 769-fold increased with anterior screw fixation, 3412-fold increased with C1-C2 posterior arthrodesis and 162-fold increased with occipito-cervical stabilization, compared with collar alone. The risk of having a worse mRS post-treatment was 11-fold increased with C1-C2 posterior arthrodesis and 15-fold increased with occipito-cervical stabilization, compared with cervical collar. Similarly, the probability of having a worse outcome measured with the SWPS increased 7 to 11-fold for patients treated with anterior screw, C1-C2 arthrodesis and occipito-cervical stabilization. Anterior odontoid screw and C1-C2 arthrodesis demonstrated identical rates (53.4%) of negative NDI (range 50-100%), and similar percentages of unsatisfactory SWPS (fair-poor), being the two treatments associated to the worst results adopting these parameters of functional outcome, although with the bias related to the crossover of those 3 patients who underwent a second step C1-C2 arthrodesis after an anterior screw mechanical failure. Occipito-cervical stabilization revealed instead unexpected positive outcomes in terms of NDI, mRS-post and SWPS respectively in 90%, 70% and 75% of cases. These data likely biased by the limited number of this subgroup of patients (n=20), all octogenarians and all died because of their severe age-related comorbidities, could be deeply influenced even by the relatively short follow-up for this subset of patients and their pre-treatment compromised general clinical status, which led these ultra-elderlies to be more accustomed to tolerate pain and to accept discomfort and functional limitations. In terms of

1 pre-injury level of function, higher comorbidity rates were associated with 10-fold increased
2 risk of worse NDI, while severe pre-treatment disabilities were associated with 14-fold
3
4 increased risk of worse SWPS.
5

6
7 Although some authors encourage the anterior odontoid screw fixation as the technique of
8
9 choice for surgical treatment of type II odontoid fractures without any substantial limitation
10
11 ^{5,6}, some others, in line with our findings recognize that morbidity, mortality and functional
12
13 outcome are influenced by patient's age ^{8,20} and age-related comorbidities ¹¹. Due to their
14
15 youthful appearance in terms of general clinical status, 11 among our ultra-elderly patients
16
17 underwent anterior odontoid screw fixation. Despite the absence of major comorbidities, this
18
19 subset of patients showed a surprisingly negative functional outcome with a 18.2% of
20
21 mechanical failure, which led to a second step C1-C2 posterior arthrodesis.
22
23
24

25
26 Therefore, differently from Henaux et al ⁴ and Hou et al ⁵, we wouldn't recommend an
27
28 anterior odontoid screw fixation in the elderly ^{4,5} whatever the fracture type, since functional
29
30 outcome appeared worse than conservative treatments, but similar to other surgical treatment
31
32 modalities (and with a theoretical lower percentage of radiological stability compared with
33
34 C1-C2 and occipito-cervical arthrodesis). Moreover, a high probability of anterior screw
35
36 loosening in these osteoporotic patients may expose both to the risk of a revision surgery
37
38 (crossover) with additional posterior fixation and to further risks in terms of morbidity and
39
40 mortality ²⁰.
41
42
43
44

45
46 In line with these trends we found a SWPS fair or poor in 53.4% of patients who underwent
47
48 anterior screw fixation, while no substantial differences in outcome were observed between
49
50 the two conservative treatments with cervical collar and halo vest, which distinguished for
51
52 excellent or good SWPS respectively in the 88.8% and 81.80% of the cases.
53
54

55
56 Joestl et al ¹⁶ emphasize the excellent or good outcome obtained in the elderly with Harms
57
58 technique up to five years after surgery. Our data, instead, showed that all surgical techniques
59
60
61
62
63
64
65

1 (anterior odontoid screw fixation, posterior arthrodesis C1-C2 with Harms technique, and
2 occipito-cervical fixation) and high mRS-pre were associated with worse results in terms of
3
4 Smiley Webster Pain Scale. Therefore, differently from Vaccaro et al ¹¹, we didn't find any
5
6 favor of surgical management in elderly patients in terms of functional outcome, but we agree
7
8 with them that patients and their families should be aware that treatment might not restore the
9
10 average patient to his/her pre-injury level of function.
11
12
13
14
15

16 **CONCLUSION**

17
18
19
20
21 Based on data validated by multiple measures of functional outcome, our study may provide
22
23 useful guidance for surgeons and clinicians in advising elderly/ultra-elderly patients and their
24
25 families about treatment for unstable odontoid fractures.
26
27

28
29 The comparison between homogenous groups of patients managed with the most adopted
30
31 conservative and surgical treatments provides a unique and comprehensive view about the
32
33 impact of each different technique on functional outcome of the most fragile patients.
34
35

36 Prospective multicenter studies evaluating clinico-radiological aspects of odontoid fractures
37
38 focused on specific types of treatment with setup based on age groups should be favoured.
39
40

41 Future studies should include measures for evaluating the level of function: radiological
42
43 outcome (fusion) should not represent the only aim of C2 fracture treatment in the elderly,
44
45 since quality of life is influenced by several factors, and a treatment modality that impairs
46
47 daily disability may have a tremendous impact also on survival of these fragile patients.
48
49

50
51 Therefore, the decision for the best treatment should be tailored taking into consideration and
52
53 balancing the probability of fusion and stability with the probability of impairing functional
54
55 outcome, the latter being negatively influenced by comorbidities (CCI), fragility (mRS) and
56
57 surgery. An analysis of these factors could lead the basis for designing algorithms able to
58
59
60
61
62
63
64
65

1 customize the treatment strategy. Further investigations, involving a larger patient sample,
2 may elucidate if the surprising advantage evidenced with conservative immobilizations,
3 especially with hard cervical collars, comes from the absence of true minimally invasive
4 procedures for odontoid fractures fixations or if materials and designs of modern cervical
5 collars can really make enough difference to represent the best treatment compromise in the
6 elderly.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES

1. Barlow DR, Higgins BT, Ozanne EM, et al. Cost Effectiveness of Operative Versus Non-Operative Treatment of Geriatric Type-II Odontoid Fracture. *Spine (Phila Pa 1976)* 2016;41:610-7.
2. Butler JS, Dolan RT, Burbridge M, et al. The long-term functional outcome of type II odontoid fractures managed non-operatively. *Eur Spine J* 2010;19:1635-42.
3. Fehlings MG, Arun R, Vaccaro AR, et al. Predictors of treatment outcomes in geriatric patients with odontoid fractures: AOSpine North America multi-centre prospective GOF study. *Spine (Phila Pa 1976)* 2013;38:881-6.
4. Henaux PL, Cueff F, Diabira S, et al. Anterior screw fixation of type IIB odontoid fractures in octogenarians. *Eur Spine J* 2012;21:335-9.
5. Hou Y, Yuan W, Wang X. Clinical evaluation of anterior screw fixation for elderly patients with type II odontoid fractures. *J Spinal Disord Tech* 2011;24:E75-81.
6. Joestl J, Lang N, Bukaty A, et al. A comparison of anterior screw fixation and halo immobilisation of type II odontoid fractures in elderly patients at increased risk from anaesthesia. *Bone Joint J* 2016;98-B:1222-6.
7. Patel A, Zakaria R, Al-Mahfoudh R, et al. Conservative management of type II and III odontoid fractures in the elderly at a regional spine centre: A prospective and retrospective cohort study. *Br J Neurosurg* 2015;29:249-53.
8. Platzner P, Thalhammer G, Ostermann R, et al. Anterior screw fixation of odontoid fractures comparing younger and elderly patients. *Spine (Phila Pa 1976)* 2007;32:1714-20.
9. Robinson Y, Robinson AL, Olerud C. Systematic review on surgical and nonsurgical treatment of type II odontoid fractures in the elderly. *Biomed Res Int* 2014;2014:231948.
10. Smith JS, Kepler CK, Kopjar B, et al. Effect of type II odontoid fracture nonunion on outcome among elderly patients treated without surgery: based on the AOSpine North America geriatric odontoid fracture study. *Spine (Phila Pa 1976)* 2013;38:2240-6.
11. Vaccaro AR, Kepler CK, Kopjar B, et al. Functional and quality-of-life outcomes in geriatric patients with type-II dens fracture. *J Bone Joint Surg Am* 2013;95:729-35.
12. Jubert P, Lonjon G, Garreau de Loubresse C. Complications of upper cervical spine trauma in elderly subjects. A systematic review of the literature. *Orthop Traumatol Surg Res* 2013;99:S301-12.

13. Molinari WJ, 3rd, Molinari RW, Khera OA, et al. Functional outcomes, morbidity, mortality, and fracture healing in 58 consecutive patients with geriatric odontoid fracture treated with cervical collar or posterior fusion. *Global Spine J* 2013;3:21-32.
14. Huybregts JG, Jacobs WC, Vleggeert-Lankamp CL. The optimal treatment of type II and III odontoid fractures in the elderly: a systematic review. *Eur Spine J* 2013;22:1-13.
15. Evaniew N, Yarascavitch B, Madden K, et al. Atlantoaxial instability in acute odontoid fractures is associated with nonunion and mortality. *Spine J* 2015;15:910-7.
16. Joestl J, Lang NW, Tiefenboeck TM, et al. Management and Outcome of Dens Fracture Nonunions in Geriatric Patients. *J Bone Joint Surg Am* 2016;98:193-8.
17. Scheyerer MJ, Zimmermann SM, Simmen HP, et al. Treatment modality in type II odontoid fractures defines the outcome in elderly patients. *BMC Surg* 2013;13:54.
18. Schroeder GD, Kepler CK, Kurd MF, et al. A Systematic Review of the Treatment of Geriatric Type II Odontoid Fractures. *Neurosurgery* 2015;77 Suppl 4:S6-14.
19. Shen Y, Miao J, Li C, et al. A meta-analysis of the fusion rate from surgical treatment for odontoid fractures: anterior odontoid screw versus posterior C1-C2 arthrodesis. *Eur Spine J* 2015;24:1649-57.
20. Lohrer L, Raschke MJ, Thiesen D, et al. Current concepts in the treatment of Anderson Type II odontoid fractures in the elderly in Germany, Austria and Switzerland. *Injury* 2012;43:462-9.

Table 1: Pre-treatment patients' characteristics

	Collar	Halo	Anterior screw	C1-C2 posterior arthrodesis	Occipito-cervical fixation	Total
<i>Age: 65-79</i>	10 (22%)	12 (54%)	19 (63%)	17 (57%)	13 (65%)	71 (48%)
<i>Age: ≥ 80</i>	35 (78%)	10 (46%)	11 (37%)	13 (43%)	7 (35%)	76 (52%)
<i>ASA: 1-2</i>	10 (23%)	9 (41%)	10 (33%)	8 (27%)	8 (40%)	45 (31%)
<i>ASA: 3</i>	33 (75%)	13 (59%)	18 (60%)	20 (67%)	10 (50%)	94 (64%)
<i>ASA: 4</i>	1 (2%)	0	2 (7%)	2 (6%)	2 (10%)	7 (5%)
<i>CCI: 2-5</i>	28 (62%)	14 (64%)	22 (73%)	23 (77%)	10 (50%)	97 (66%)
<i>CCI: 6-8</i>	17 (38%)	8 (36%)	8 (27%)	7 (33%)	10 (50%)	50 (34%)
<i>mRS-pre: 0-3</i>	35 (78%)	21 (95%)	14 (47%)	13 (43%)	20 (100%)	103 (70%)
<i>mRS-pre: 4-5</i>	10 (22%)	1 (5%)	16 (43%)	17 (47%)	0	44 (30%)
<i>Dislocation < 5 mm</i>	44 (97%)	21 (91%)	29 (96%)	26 (87%)	14 (72%)	135 (92%)
<i>Dislocation ≥ 5 mm</i>	1 (3%)	1 (9%)	1 (4%)	4 (13%)	6 (28%)	12 (8%)

ASA, ASA score; CCI, Charlson Comorbidity Index; mRS-pre, pre-treatment modified Rankin Scale.

Table 2: Analysis of factors affecting outcome in relation to NDI (0-48 Vs 50-100)

Logistic regression analysis: dichotomous NDI	p	OR	95% C.I. for OR Lower	95% C.I. for OR Upper
<i>Gender (male Vs female)</i>	0,330	2,375	0,417	13,524
<i>Age ≥ 80</i>	0,956	0,947	0,136	6,583
<i>CCI: 6 or 7 or 8</i>	0,039	10,260	1,127	93,384
<i>ASA 2</i>	0,130			
<i>ASA 3</i>	0,064	8,067	0,887	73,327
<i>ASA 4</i>	0,815	0,530	0,003	108,380
<i>mRS-pre: 4 or 5</i>	0,001	159,676	9,037	2821,409
<i>Anderson D'Alonzo Classification 2</i>	0,409			
<i>Anderson D'Alonzo Classification 3</i>	0,662	1,815	0,126	26,200
<i>Anderson D'Alonzo Classification 2 + other fractures of cranio-vertebral junction</i>	0,188	3,525	0,541	22,982
<i>Dislocation ≥ 5mm</i>	0,244	0,052	0,0001	7,492
<i>Treatments: cervical collar</i>	0,005			
<i>halo vest</i>	0,035	89,109	1,373	5783,955
<i>odontoid anterior screw fixation</i>	0,001	769,807	17,959	32997,711
<i>posterior arthrodesis (Harms)</i>	0,0001	3412,629	44,766	260153,092
<i>occipito-cervical stabilization</i>	0,013	162,042	2,899	9058,388

ASA, ASA score; CCI, Charlson Comorbidity Index; mRS-pre, pre-treatment modified Rankin Scale.

Table 3: Analysis of factors affecting outcome in relation to SWPS

Logistic regression analysis: SWPS	p	OR	95% C.I. for OR Lower	95% C.I. for OR Upper
<i>Gender (male Vs female)</i>	0,233	1,899	0,662	5,449
<i>Age ≥ 80</i>	0,959	0,968	0,277	3,378
<i>CCI: 6 or 7 or 8</i>	0,662	0,772	0,242	2,462
<i>ASA 2</i>	0,448			
<i>ASA 3</i>	0,206	2,106	0,664	6,674
<i>ASA 4</i>	0,556	2,048	0,189	22,243
<i>mRS-pre: 4 or 5</i>	0,0001	14,394	3,732	55,512
<i>Anderson D'Alonzo Classification 2</i>	0,334			
<i>Anderson D'Alonzo Classification 3</i>	0,336	0,432	0,078	2,390
<i>Anderson D'Alonzo Classification 2 + other fractures of cranio-vertebral junction</i>	0,183	0,419	0,116	1,509
<i>Dislocation ≥ 5mm</i>	0,329	0,342	0,040	2,946
<i>Treatments: cervical collar</i>	0,047			
<i>halo vest</i>	0,040	6,409	1,086	37,820
<i>odontoid anterior screw fixation</i>	0,004	11,523	2,221	59,787
<i>posterior arthrodesis (Harms)</i>	0,022	7,030	1,321	37,405
<i>occipito-cervical stabilization</i>	0,016	11,355	1,568	82,243

ASA, ASA score; CCI, Charlson Comorbidity Index; mRS-pre, pre-treatment modified Rankin Scale.

Table 4: Analysis of factors affecting outcome in relation to mRS-post

Logistic regression analysis: mRS-post	p	OR	95% C.I. for OR Lower	95% C.I. for OR Upper
<i>Gender (male Vs female)</i>	0,831	1,116	0,410	3,037
<i>Age ≥ 80</i>	0,086	2,820	0,864	9,201
<i>CCI: 6 or 7 or 8</i>	0,219	1,982	0,666	5,899
<i>ASA 2</i>	0,788			
<i>ASA 3</i>	0,953	0,966	0,306	3,051
<i>ASA 4</i>	0,521	0,479	0,050	4,551
<i>mRS-pre: 4 or 5</i>	0,185	2,324	0,668	8,088
<i>Anderson D'Alonzo Classification 2</i>	0,886			
<i>Anderson D'Alonzo Classification 3</i>	0,640	1,403	0,340	5,793
<i>Anderson D'Alonzo Classification 2 + other fractures of cranio-vertebral junction</i>	0,786	1,186	0,345	4,078
<i>Dislocation ≥ 5mm</i>	0,208	2,739	0,570	13,153
<i>Treatments: cervical collar</i>	0,023			
<i>halo vest</i>	0,142	4,031	0,627	25,911
<i>odontoid anterior screw fixation</i>	0,141	3,669	0,649	20,732
<i>posterior arthrodesis (Harms)</i>	0,004	11,092	2,163	56,874
<i>occipito-cervical stabilization</i>	0,005	15,770	2,275	109,343

ASA, ASA score; CCI, Charlson Comorbidity Index; mRS-pre, pre-treatment modified Rankin Scale.

Table 5: Analysis of factors affecting outcome (all patients): risk of crossover

Logistic regression analysis: risk of crossover	p	OR	95% C.I. for OR Lower	95% C.I. for OR Upper
<i>Gender (male Vs female)</i>	0,512	0,720	0,270	1,919
<i>Age ≥ 80</i>	0,004	0,180	0,056	0,577
<i>CCI: 6 or 7 or 8</i>	0,424	0,624	0,197	1,980
<i>ASA 2</i>	0,542			
<i>ASA 3</i>	0,347	1,692	0,565	5,066
<i>ASA 4</i>	0,396	3,779	0,175	81,499
<i>mRS-pre: 4 or 5</i>	0,480	1,662	0,405	6,813
<i>Anderson D'Alonzo Classification 2</i>	0,478			
<i>Anderson D'Alonzo Classification 3</i>	0,314	0,310	0,032	3,034
<i>Anderson D'Alonzo Classification 2 + other fractures of cranio-vertebral junction</i>	0,610	1,327	0,447	3,938
<i>Dislocation ≥ 5mm</i>	0,331	0,322	0,033	3,158
<i>Treatments: cervical collar</i>	0,009			
<i>halo vest</i>	0,255	1,963	0,614	6,271
<i>odontoid anterior screw fixation</i>	0,056	0,225	0,049	1,041
<i>posterior arthrodesis (Harms)</i>	0,035	0,090	0,010	0,843
<i>occipito-cervical stabilization</i>	0,125	0,171	0,018	1,632

ASA, ASA score; CCI, Charlson Comorbidity Index; mRS-pre, pre-treatment modified Rankin Scale.

Figure Legend:

Figure 1: NDi distribution in the different treatments

Figure 1: NDI distribution in the different treatments

