

This is a pre print version of the following article:

Electrophysiological characterization and functionality of neurons in murine cerebral organoids / Curia, G., Ren, E., Bardoni, R., Decimo, I., Ciarpella, F., Zamfir, R.g., Campanelli, A.. - (2021).
[10.5281/zenodo.7600194]

Zenodo

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

26/10/2026 19:07

(Article begins on next page)

Electrophysiological characterization and functionality of neurons in murine cerebral organoids

doi: [10.5281/zenodo.7600194](https://doi.org/10.5281/zenodo.7600194)

GENERAL INFORMATION

- Organoids: Brain organoids were obtained starting from E14.5 murine-derived Neural Stem Cells (NSCs). Organoids at different stages of development (11-13 DIV: intermediate; 19-26 DIV: early mature; 33-38 DIV: mature) have been used.
- Extracellular solution, artificial cerebrospinal fluid (ACSF), (in mM): NaCl (115.0), KCl (2.0), KH_2PO_4 (1.25), MgSO_4 (1.0), CaCl_2 (2.0), D-Glucose (25.0), and NaHCO_3 (26.0), and bubbled in carbogen (95% O_2 and 5% CO_2). When indicated, tetrodotoxin (TTX) applied was 1 μM (protocols 0003 and 0004).
- Intracellular solution (in mM): K-Gluconate (120.0), NaCl (5), HEPES (10.0), EGTA (10.0), MgCl_2 (2.0), CaCl_2 (1.0), ATP- Na_2 (2.0), and GTP- Na (0.4), osmolality was adjusted at 285 mOsm with 10 mM sucrose.
- Acquisition system: data were acquired in whole-cell configuration with an Axopatch 1D amplifier and Digidata 1322, using pClamp 10.6 software (all from Molecular Devices, USA).
- For voltage-clamp (VC) recordings, membrane potential was held at -70 mV and positive voltage steps (500 ms) were applied from -70 mV to +100 mV ($\Delta V=10$ mV).
- Current-clamp (CC) activity was recorded at resting membrane potential and evoked by 500 ms current pulses from -10 pA to +10 pA ($\Delta A=10$ pA).

DATASET FORMAT

The experiments folders are compressed in *.zip format.

Below summarized the file formats and the software with which they can be opened:

File format	Required software
.abf	pClamp or Clampfit software (Molecular Devices, USA)

DATASET STRUCTURE

- main folder [Organoidi XX-XX DIV]
 - Organoid folder [Organoid XX DIV_DD.MM.YYYY]
 - *.abf recording files including voltage-clamp (VC) and current-clamp (CC) protocols:
 - 0000 VC step in ACSF
 - 0001 CC step in ACSF
 - 0002 CC step in ACSF at Rheo base
 - 0003 VC step in ACSF+TTX
 - 0004 CC step in ACSF+TTX

FILE NAMING STRUCTURE

Date [YYYYMMDD]_Cell number [CellXX]_Organoid number [OrganoidXX]_Protocol [XXXX]

- e.g.: 20191217_Cell03_Organoid01_0000.abf: experiment performed on December 17th 2019, cell number 3 recorded from organoid number 1, protocol VC step in ACSF.