

References

- Accelrys Software Inc., *Discovery Studio Modeling Environment, Release 3.5, San Diego: Accelrys Software Inc., 2012.*
- Amiri, S., Tai, K., Beckstein, O., Biggin, P.C., Sansom, M.S.P. (2005) The $\alpha 7$ nicotinic acetylcholine receptor: molecular modelling, electrostatics, and energetic. *Mol. Membrane Biol.* 22(3), 151-162.
- Baker, N.A., Sept, D., Joseph, S., Holst, M.J., and McCammon, J.A. (2001). Electrostatics of nanosystems: application to microtubules and the ribosome. *PNAS, USA* 98, 10037-10041.
- Barnes, M.N., Hales, T.G., Lummis, S.C.R. and Peters, J.A. (2009) The 5-HT₃ receptor – the relationship between structure and function. *Neuropharm.* 56, 273-284.
- Barrera, N. P., Herbert, P., Henderson, R. M., Martin, I. L. and Edwardson, J. M. (2005) Atomic force microscopy reveals the stoichiometry and subunit arrangement of 5-HT₃ receptors. *PNAS.* 102, 12595-12600.
- Barry, P.H., and Lynch, J.W. (2005). Ligand-Gated Channels. *IEEE Trans. Nanobio.,* 4, 71-80.
- Beckstein, O., Tai, K., & Sansom, M.S.P. (2004). Not ions alone: barriers to ion permeation in nanopores and channels. *J. Am. Chem. Soc.* 126, 14694–14695.
- Beckstein, O., and Sansom, M.S.P. (2006). A hydrophobic gate in an ion channel: the closed state of the nicotinic acetylcholine receptor. *Phys. Biol.,* 3, 147-159.
- Beene, D.L., Brandt, G.S., Zhong, W., Zacharias, N.M., Lester, H.A., Dougherty, D.A. (2002) Cation- π Interactions in Ligand Recognition by Serotonergic (5-HT_{3A}) and Nicotinic Acetylcholine Receptors: The Anomalous Binding Properties of Nicotine. *Biochemistry,* 41, 10262–10269.
- Berendsen, H. J. C., van der Spoel, D., and van Drunen, R. (1995) GROMACS: a message-passing parallel molecular dynamics implementation. *Comput. Phys. Commun.,* 91, 43-56.
- Bieber, T., and Elsässer, H.P. (2001) Preparation of a low molecular weight polyethylenimine for efficient cell transfection. *Biotechniques.* 30(1), 74-7, 80-1.
- Boess, F.G., Lummis, S.C.R. and Martin, I.L. (1992) Molecular Properties of 5-Hydroxytryptamine₃ Receptor-Type Binding Sites Purified from NG108-15 Cells. *J. Neurochem.* 59, 1692-1701.

- Boess, F.G., Beroukhim, R. and Martin, I.L. (1995) Ultrastructure of the 5-hydroxytryptamine₃ receptor. *J. Neurochem.* 64, 1401-05.
- Boess, F.G., Steward, L.J., Steele, J.A., Liu, D., Reid, J., Glencorse, T.A., and Martin, I.L. (1997) Analysis of the Ligand Binding Site of the 5-HT₃ Receptor Using Site Directed Mutagenesis: Importance of Glutamate 106. *Neuropharmacology*, **36**, 637-647
- Bocquet, N., Nury, H., Baaden, M., Le Poupon, C., Changeux, J. P., Delarue, M. and Corringer, P. J. (2009). X-ray structure of a pentameric ligand-gated ion channel in an apparently open conformation. *Nature*, *457*, 111-114.
- Bouzat, C. (2012) New insights into the structural bases of activation of Cys-loop receptors. *J. Pharmacol.* 106, 23-33.
- Boyd, G.W., Low, P., Dunlop, J.I., Robertson, L.A., Vardy, A., Lambert, J.L., Peters, J.A., Connolly, C.N. (2002) Assembly and Cell Surface Expression of Homomeric and Heteromeric 5-HT₃ Receptors: The Role of Oligomerization and Chaperone Proteins. *Mol. Cell. Neurosci.* 21, 38–50.
- Brady, C.A., Stanford, I.M., Ali, I., Lin, L., Williams, J.M., Dubin, A.E., Hope, A.G., Barnes, N.M. (2001) Pharmacological comparison of human homomeric 5-HT_{3A} receptors versus heteromeric 5-HT_{3A/3B} receptors. *Neuropharm.* 41, 282–284.
- Brejck, K., van Dijk, W.J., Klaassen, R.V., Schuurmans, M., van Der Oost, J., Smit, A.B. and Sixma, T.K. (2001) Crystal structure of an ACh-binding protein reveals the ligand-binding domain of nicotinic receptors. *Nature*, 411, 269–276.
- Buchan, D.W., Ward, S.M., Loble, A.E., Nugent, T.C., Bryson, K., and Jones, D.T. (2010). Protein annotation and modelling servers at University College London. *Nucleic Acids Res.*, 38 Suppl, W563-W568.
- Butler, A., Hill, J.M., Ireland, S.J., Jordan, C.C., and Tyers, M.B. (1988) Pharmacological properties of GR38032F, a novel antagonist at 5-HT₃ receptors. *Br. J. Pharmacol.* 94, 397-412.
- Cappelli, A., Anzini, M., Vomero, S., Mennuni, L., Makovec, F., Doucet, E., Hamon, M., Bruni, G., Romeo, M.R., Menziani, M.C., De Benedetti, P.G., and Langer, T. (1998) Novel Potent and Selective Central 5-HT₃ Receptor Ligands Provided with Different Intrinsic Efficacy. 1. Mapping the Central 5-HT₃ Receptor Binding Site by Arylpiperazine Derivatives. *J. Med. Chem.*, 41, 728.

- Cappelli, A., Anzini, M., Vomero, S., Mennuni, L., Makovec, F., Hamon, M., De Benedetti, P.G., and Menziani, M.C. (2002) The Interactions of the 5-HT₃ receptor with arylpiperazine, tropane, and quinuclidine ligands. *Curr. Top. Med. Chem.* 2, 599-624.
- Cappelli, A., Gallelli, A., Manini, M., Anzini, M., Mennuni, L., Makovec, F., Menziani, M.C., Alcaro, S., Ortuso, F., and Vomero, S. (2005) Further studies on the interaction of the 5-hydroxytryptamine₃ (5-HT₃) receptor with arylpiperazine ligands. development of a new 5-HT₃ receptor ligand showing potent acetylcholinesterase inhibitory properties. *J. Med. Chem.* 48, 3564-3575.
- Cappelli, A., Butini, S., Brizzi, A., Gemma, S., Valenti, S., Giuliani, G., Anzini, M., Mennuni, L., Campiani, G., Brizzi, V., and Vomero, S. (2010) The interactions of the 5-HT₃ receptor with quipazine-like arylpiperazine ligands: the journey track at the end of the first decade of the third millennium. *Curr. Top. Med. Chem.* 10, 504-526.
- Clark, R.D., Miller, A.B., Berger, J., Repke, D.B., Weinhardt, K.K., Kowalczyk, B.A., Eglén, R.M., Bonhaus, D.W., Lee, C.H., Michel, A.D., Smith, W.L., Wong, E.H.F. (1993) 2-(Quinuclidin-3-yl)pyrido[4,3-b]indol-1-ones and isoquinolin-1-ones. Potent conformationally restricted 5-HT₃ receptor antagonists. *J. Med.Chem.*,36, 2645.
- Colovos, C., and Yates, T.O. (1993). Verification of protein structures: patterns of nonbonded atomic interactions. *Prot. Science*, 2(9), 1511-9.
- Corringer, P.J., Bertrand, S., Galzi, J.L., Devillers-Thiery, A., Changeux, J.P., Bertrand, D. (1999) Mutational analysis of the charge selectivity filter of the alpha7 nicotinic acetylcholine receptor. *Neuron*. 22, 831-843.
- Corringer, P.J., Le Novère, N., and Changeux, J. P. (2000). Nicotinic receptors at the amino acid level. *Annu. Rev. Pharmacol. Toxicol.*, 40, 431-458.
- Cymes, G.D., and Grosman, C. (2011) Tunable pK_a values and the basis of opposite charge selectivities in nicotinic-type receptors. *Nature*, 474, 526-530.
- Dang, H., England, P.M., Farivar, S.S., Dougherty, D.A., and Lester, H.A. (2000). Probing the role of a conserved M1 proline residue in 5-hydroxytryptamine₃ receptor gating. *Mol. Pharm.*, 57, 1114-1122.
- Davies, P.A., Pistis, M., Hanna, M.C., Peters, J.A., Lambert, J.J., Hales, T.G., Kirkness, E.F. (1999) The 5-HT_{3B} subunit is a major determinant of serotonin-receptor function. *Nature* 397, 359-363.

- De Rienzo, F., Moura Barbosa, A.J., Perez, M.A.S., Fernandes, P.A., Ramos, M.J., and Menziani, M.C. (2012) The extracellular subunit interface of the 5-HT₃ receptors: a computational alanine scanning mutagenesis study. *J. Biomol. Struct.Dyn.* 30, 280-298.
- DeLano, W. L. (2008) PyMOL Molecular Graphics System, *DeLano Scientific LLC*, Palo Alto, CA.
- Dolinsky, T.J., Nielsen, J.E., McCammon, J.A., and Baker, N.A. (2004). PDB2PQR: an automated pipeline for the setup, execution, and analysis of Poisson-Boltzmann electrostatics calculations. *Nucleic Acids Res.* 32, W665-W667.
- Dolinsky, T.J., Czodrowski, P., Li, H., Nielsen, J.E., Jensen, J.H., Klebe, G., & Baker, N.A. (2007). PDB2PQR: Expanding and upgrading automated preparation of biomolecular structures for molecular simulations. *Nucleic Acids Res.* 35, W522-5.
- Dubin, A.E., Huvar, R., D'Andrea, M.R., Pyati, J., Zhu, J.Y., Joy, K.C., Wilson, S.J., Galindo, J.E., Glass, C.A., Luo, L., Jackson, M.R., Lovenberg, T.W., Erlander, M.G. (1999) The Pharmacological and Functional Characteristics of the Serotonin 5-HT_{3A} Receptor Are Specifically Modified by a 5-HT_{3B} Receptor Subunit. *J. Bio. Chem.* 274(43), 30799–30810.
- Eisemberg, D., Weiss, R.M., Terwilliger, T.C., and Wilcox, W. (1982) Hydrophobic moments and protein structure. *Faraday Symp. Chem. Soc.* 17, 109-120.
- Essmann, U., Perera, L., Berkowitz, M.L., Daren, T., Lee, H., and Pedersen, L.G. (1995) A smooth particle mesh Ewald method. *J. Chem. Phys.*, 103, 8577-8593.
- Filatov, G.N., and White, M.M. (1995). The Role of Conserved Leucines in the M2 Domain of the Acetylcholine Receptor in Channel Gating. *Mol. Pharm.*, 48, 379-384.
- Fozard, J.R. and Host, M., (1982) Selective inhibition of the Bezold-Jarish effect of 5-HT in the rat by antagonists at neuronal 5-HT receptors. *Br. J. Pharmac.* 57, 115-125.
- Fozard, J.R. (1984) MDL 72222: a potent and highly selective antagonist at neuronal 5-hydroxytryptamine receptors. *Naunyn Schmiedebergs Arch. Pharmacol.* 326, 36-44.
- Funahashi, M., Mitoh, Y. and Matsuo, R. (2004) Activation of presynaptic 5-HT₃ receptors facilitates glutamatergic synaptic inputs to area postrema neurons in rat brain slices. *Methods Find Exp. Clin. Pharmacol.* 26, 615-22.
- Gabdoulline, R.R, Wade, R.C., and Walther, D. (1999) MolSurfer: two dimensional maps for navigating three-dimensional structures of proteins. *Trends Biochem. Sci.* 24, 285-287.

- Gabdoulline, R.R, Wade, R.C., and Walther, D. (2003) MolSurfer: a macromolecular interface navigator. *Nucleic. Acids Res.* 31, 3349-3351.
- Gaddum, J.H. and Picarelli, Z.P. (1957) Two kinds of tryptamine receptor. *Brit. J. Pharmacol.* 12, 323.
- Galeazzi, S., Hermans, T.M., Paolino, M., Anzini, M., Mennuni, L., Giordani, A., Caselli, G., Makovec, F., Meijer, E.W., Vomero, S., and Cappelli, A. (2010) Multivalent Supramolecular Dendrimer-Based Drugs. *Biomacromol.*, 11, 182-186.
- Glennon, R.A. (2006) Strategies for the Development of Selective Serotonergic Agents. *The receptors* 91-142.
- Greenshaw, A.J. and Silverstone, P.H. (1997) The non-antiemetic uses of serotonin 5-HT₃ receptor antagonists. Clinical pharmacology and therapeutic applications. *Drugs.* 53, 20-39.
- Grosman, C., Zhou, M., and Auerbach, A. (2000). Mapping the conformational wave of acetylcholine receptor channel gating. *Nature*, 403, 773-776.
- Gunthorpe, M.J., and Lummis, S.C.R. (2001). Conversion of the ion selectivity of the 5-HT_{3A} receptor from cationic to anionic reveals a conserved feature of the ligand gated ion channel superfamily. *J. Bio. Chem.*, 276, 10977-10983.
- Hales, T.G., Dunlop, J.I., Deeb, T.Z., Carland, J.E., Kelley, S.P., Lambert, J.J., and Peters, J.A. (2006). Common determinants of single channel conductance within the large cytoplasmic loop of 5-hydroxytryptamine type 3 and $\alpha 4\beta 2$ nicotinic acetylcholine receptors. *J. Bio. Chem.*, 281, 8062-8071.
- Hanna, M.C., Davies, P.A., Hales, T.G., Kirkness, E.F. (2000) Evidence for Expression of Heteromeric Serotonin 5-HT₃ receptors in Rodents. *J. Neurochem.* 75, 240-247.
- Hannon, J. and Hoyer, D. (2008) Molecular biology of 5-HT receptors. *Behav Brain Res.* 195, 198-213.
- Hapfelmeier, G., Tredt, C., Haseneder, R., Zieglgansberger, W., Eisensamer, B., Rupprecht, R., Rammes, G. (2003) Co-expression of the 5-HT_{3B} Serotonin Receptor Subunit Alters the Biophysics of the 5-HT₃ Receptor. *Biophysic. J.* 84, 1720–1733.
- Heidempergher, F., Pillan, A., Pinciroli, V., Vaghi, F., Arrigoni, C., Bolis, G., Caccia, C., Dho, L., McArthur, R., and Varasi, M. (1997) Phenylimidazolidin-2-one Derivatives as Selective 5-HT₃ Receptor Antagonists and Refinement of the Pharmacophore Model for 5-HT₃ Receptor Binding. *J. Med. Chem.*, 40, 3369.

- Herrstedt, J., Dombernowsky, P. (2007) Anti-emetic therapy in cancer chemotherapy: current status. *Basic Clin. Pharmacol. Toxicol.* 101, 143-50.
- Hess, B.; Kutzner, C., van der Spoel, D., and Lindahl, E. (2008) GROMACS 4: Algorithms for Highly Efficient, Load-Balanced, and Scalable Molecular Simulation. *J. Chem. Theory Comput.*, 4 (3), 435–447 .
- Hibbs, R.E. and Gouaux, E. (2011). Principles of activation and permeation in an ion-selective Cys-loop receptor. *Nature*, 474, 54-60.
- Hilf, R.J.C. and Dutzler, R. (2008) X-ray structure of a prokaryotic pentameric ligand-gated ion channel. *Nature*, 452, 375-380.
- Holbrook, J. D., Gill, C. H., Zebda, N., Spencer J. P., Leyland, R., Rance, K. H., Trin, H., Balmer, G., Kelly, F. M., Yusaf, S. P., Courtenay, N., Luck, J., Rhodes, A., Modha, S., Moore, S. E., Sanger, G. J. and Gunthorpe, M. J. (2009) Characterisation of 5-HT_{3C}, 5-HT_{3D} and 5-HT_{3E} receptor subunits: evolution, distribution and function. *J. Neurochem.* 108, 384-396.
- Hooft, R.W.W., Vriend, G., Sander, C., and Abola, E.E. (1996). Errors in protein structures, *Nature*, 381, 272-272.
- Humphrey, W., Dalke, A., and Schulten, K. (1996) VMD - Visual Molecular Dynamics, *J. Molec. Graphics*, 14, 33-38.
- Hung, A., Tai, K., and Sansom, M.S.P. (2005). Molecular Dynamics Simulation of the M2 helices within the Nicotinic Acetylcholine Receptor Transmembrane Domain: Structure and Collective Motions. *Biophysic Journal*, 88, 3321-3333.
- Hussy, N., Lukas, W. and Jones, K.A. (1994) Functional properties of a cloned 5-hydroxytryptamine ionotropic receptor subunit: comparison with native mouse receptors. *J. Physiol.* 481,2.
- Jensen, A.A., Davies, P.A., Brauner-Osborne, H., and Krzywkowski, K. (2008) 3B but which 3B? And that's just one of the questions: the heterogeneity of human 5-HT₃ receptors. *Trends in Pharmacol Sci*, 29, 437-4
- Jones, D.T. (1999). Protein secondary structure prediction based on position-specific scoring matrices. *J. Mol. Bio.*, 292, 195-202.
- Joshi, P.R., Suryanarayanan, A., Hazai, E., Schulte, M.K., Maksay, G. and Bikádi, Z. (2006) Interactions of Granisetron with an agonist-free 5-HT_{3A} receptor model. *Biochem.* 45, 1099–1105.

- Kaihsu, T., Fowler, P., Mokrab, Y., Stansfeld, P., & Sansom, S.P. (2008). Molecular Modeling and Simulation Studies of Ion Channel Structures, Dynamics and Mechanisms. *Meth. Cell Bio.* 90(12), 233-265.
- Karlin, A. (2002) Emerging structure of the nicotinic acetylcholine receptors. *Nat. Rev. Neurosci.* 3, 102-114.
- Karnovsky, A.M., Gotow, L.F., McKinley, D.D., Piechan, J.L., Rublea, C.L., Mills, C.J., Schellin, K.A.B., Slightom, J.L., Fitzgerald, L.R., Benjamin, C.W. and Roberds, S.L. (2003) A cluster of novel serotonin receptor 3-like genes on human chromosome 3. *Gene* 319, 137–148.
- Kelley, S.P., Dunlop, J.I., Kirkness, E.F., Lambert, J.J., and Peters, J.A. (2003). A cytoplasmic region determines single-channel conductance in 5-HT₃ receptors. *Nature*, 424, 321-324.
- Kienker, P., Tomaselli, G., Jurman, M., and Yellen, G. (1994). Conductance mutations of the nicotinic acetylcholine receptor do not act by a simple electrostatic mechanism. *Biophysic. J.*, 66, 325–334.
- Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E.L. (2001). Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Bio.*, 305(3), 567-580.
- Kunkel, T.A. (1985) Rapid and efficient site-specific mutagenesis without phenotypic selection. *Proc. Natl. Acad. Sci. USA* 82(2), 488-92.
- Kuyucak, S., Hoyles, M., Chung, S. (1998) Analytical Solutions of Poisson's Equation for Realistic Geometrical Shapes of Membrane Ion Channels. *Biophys. J.* 74(1), 22-36.
- Laskowski, R.A., MacArthur, M.W., Moss, D.S., and Thornton, J.M. (1993). PROCHECK: a program to check the stereochemical quality of protein structure, *J. App. Cryst.*, 26, 283-291.
- Le, N. N. and Changeux, J. P. (2001) LGICdb: the ligand-gated ion channel database. *Nucleic Acids Res.* 29, 294-295.
- Lerner, M.G., and Carlson, H.A. (2006). APBS plugin for PyMOL, University of Michigan, Ann Arbor.
- Lester, H.A., Dibas, M.I., Dahan, D.S., Leite, J.F., and Dougherty, D.A. (2004) Cys-loop receptors: new twists and turns. *Trends Neurosci.* 27, 329-336.

- Liu, X., Xu, Y., Li, H., Wang, X., Jiang, H., and Barrantes, F.J. (2008). Mechanics of Channel Gating of the Nicotinic Acetylcholine Receptor. *PLoS Comp. Biol.*, 4, 100-110.
- Livesey, M.R., Cooper, M.A., Deep, T.Z., Carland, J.E., Kozuska, J., Hales, T.G., Lambert, J.J., and Peters, J.A. (2008). Structural determinants of Ca²⁺ permeability and conduction in the human 5-HT₃A receptor. *J. Bio. Chem.*, 282, 6172-6182.
- Lochner, M. and Lummis, S.C.R. (2010) Agonists and antagonists bind to an A-A interface in the heteromeric 5-HT₃AB receptor. *Biophys. J.* 98(8), 1494-502.
- Lummis, S.C.R. (2012) 5-HT₃ receptors. *J. Biol. Chem.* 287(48), 40239-45.
- Machu, T.K. (2011) Therapeutics of 5-HT₃ receptor antagonists: Current uses and future directions. *Pharmacol. Therap.* 130, 338-347.
- Madura, J.D., Briggs, J.M., Wade, R.C., Davis, R., Luty, B.A., Ilin, A., Antonsiewicz, J., Bagheri, M.K., Scott, B., and McCammon, J.A. (1995) Electrostatics and diffusion of molecules in solution: simulations with the University of Houston Brownian Dynamics program. *Comp. Phys. Comm.* 91, 57-95.
- Maksay, G., Bikadi, Z. and Simonyi, M. (2003) Binding interactions of antagonists with 5-hydroxytryptamine 3A receptor models. *J. Recept. Signal Transduct. Res.* 23, 255-270.
- Maricq, A.V., Peterson, A.S., Brake, A.J., Myers, R.M. and Julius, D. (1991) Primary structure and functional expression of the 5-HT₃ receptor, a serotonin gated-ion channel. *Science* 254, 432-437.
- Menziani, M.C., De Rienzo, F., Cappelli, A., Anzini, M. and De Benedetti, P.G. (2001) A computational model of the 5-HT₃ receptor extracellular domain: search for ligand binding sites. *Theor. Chem. Acc.* 106, 98-104.
- Miller, C. (1989). Genetic manipulation of ion channels: a new approach to structure and mechanism. *Neuron.*, 2, 1195-1205.
- Miyazawa, A., Fujiyoshi, Y. and Unwin, N. (2003) Structure and gating mechanism of the acetylcholine receptor pore. *Nature*, 423, 949-955.
- Mokrab, Y., Bavro, V.N., Mizuguchi, K., Todorov, N.P., Martin, I.L., Dunn, S.M.J., Chan, S.L., Chau, P.L. (2007) Exploring ligand recognition and ion flow in comparative models of the human GABA type A receptor. *J. Mol. Graph. Model.* 26, 760-774.
- Morris, G.M., Goodsell, D.S., Halliday, R.S., Huey, R., Belew, R.K., and Olson, A.J. (1998) Automated Docking Using a Lamarckian Genetic Algorithm and Empirical Binding Free Energy Function, *J. Comp Chem.*, 19, 1639-1662.

- Moura Barbosa, A.J., De Rienzo, F., Ramos, M.J. and Menziani, M.C. (2010) Computational analysis of ligand recognition sites of homo- and heteropentameric 5-HT₃ receptors. *Eur. J. Med. Chem.* 45(11),4746-60.
- Nelson, D.R., and Thomas, D.R. (1989) [³H]-BRL 43694 (Granisetron), a Specific Ligand for 5-HT₃ Binding Sites in Rat Brain Cortical Membranes. *Biochem. Pharmacol.*, 38, 1693-1695.
- Niesler, B., Frank, B., Kapeller, J. and Rappold, G.A. (2003) Cloning, physical mapping and expression analysis of the human 5-HT₃ serotonin receptor-like genes HTR3C, HTR3D and HTR3E. *Gene* 22, 310, 101-11.
- Niesler, B., Walstab, J., Combrink, S., Moller, D., Kapeller, J., Rietdorf, J., Bonisch, H., Gothert, M., Rappold, G. and Bruss, M. (2007) Characterization of the Novel Human Serotonin Receptor Subunits 5-HT_{3C}, 5-HT_{3D} and 5-HT_{3E}. *Mol. Pharmacol.* 72, 8-17.
- Ortells, M.O., and Barrantes, G.E. (2008) A model for the assembly of nicotinic receptors based on subunit-subunit interactions. *Proteins*, 70, 473-488.
- Paniker, S., Cruz, H., Arrabit, C., and Slesinger, A. (2002). Evidence for a centrally located gate in the pore of a serotonin-gated ion channel. *J. Neurosci.*, 22, 1629-1639.
- Parsegian, A. (1969) Energy of an Ion crossing a Low Dielectric Membrane: Solutions to Four Relevant Electrostatic Problems. *Nature*, 221, 844 - 846.
- Partenskij, M.B., Jordan, P.C. (1992) Theoretical perspectives on ion-channel electrostatics: continuum and microscopic approaches. *Q. Rev. Biophys.* 25, 477-510.
- Peters, J.A., Hales, T.G. and Lambert, J.J. (2005) Molecular determinants of single-channel conductance and ion selectivity in the Cys-loop family: insights from the 5-HT₃ receptor. *Trends Pharmacol. Sci.* 26(11), 587-594.
- Peters, J.A., Cooper, M.A., Carland, J.E., Livesey, M.R., Hales, T.G., and Lambert, J.J. (2010). Novel structural determinants of single channel conductance and ion selectivity in 5-hydroxytryptamine type 3 and nicotinic acetylcholine receptors. *J. Phys.*, 588(4), 587-595.
- Price, K.L. Lummis, S.C.R. (2004) The Role of Tyrosine Residues in the Extracellular Domain of the 5-Hydroxytryptamine₃ Receptor *J. Biol. Chem.*, 279, 23294-23301.
- Price, K.L., Bower, K.S., Thompson, A.J., Lester, H.A., Dougherty, D.A. and Lummis, S.C.R. (2008) A Hydrogen Bond in Loop A Is Critical for the Binding and Function of the 5-HT₃ Receptor. *Biochem.*, 47, 6370-6377.

- Rapport, M.M, Green, A.A., and Page, I.H. (1947) Purification of the substance which is responsible for the vasoconstrictor activity of serum. *Fed. Proc.* 6, 184.
- Rashin, A., Hoing, B. (1985) Reevaluation of the Born model of ion hydration. *J. Phys. Chem.* 89, 5588.
- Reeves, D.C., Lummis, S.C.R. (2002) The molecular basis of the structure and function of the 5-HT₃ receptor – a model ligand-gated ion channel. *Mol. Membr. Biol.* 19, 11-26.
- Reeves, D.C., Sayed, M.F., Chau, P.L., Price, K.L. and Lummis, S.C.R. (2003) Prediction of 5-HT₃ receptor agonist-binding residues using homology modeling. *Biophys. J.* 84, 2338–2344.
- Richardson, B.P., Engel, G., Donatsch, P. and Stadler, P.A. (1985) Identification of serotonin M-receptor subtypes and their specific blockade by a new class of drugs. *Nature* 316, 126-131.
- Rojas, C., Stathis, M., Thomas, A.G., Massuda, E.B., Alt, J., Zhang, J., Rubenstein, E., Sebastiani, S., Cantoreggi, S., Snyder, S.H., Slusher, B. (2008) Palonosetron Exhibits Unique Molecular Interactions with the 5-HT₃ Receptor. *Anesth Analg.* 107, 469-78.
- Rojas, C., Li, Y., Zhang, J., Stathis, M., Alt, J., Thomas, A.G., Cantoreggi, S., Sebastiani, S., Pietra, C., and Slusher, B.S. (2010) The antiemetic 5-HT₃ receptor antagonist Palonosetron inhibits substance P-mediated responses in vitro and in vivo. *J. Pharmacol. Exp. Ther.*, 335(2), 362-8.
- Rondé, P. and Nichols, R.A. (1998) High calcium permeability of serotonin 5-HT₃ receptors on presynaptic nerve terminals from rat striatum. *J. Neurochem.* 70, 1094-103.
- Sali, A., and Blundell, T.L. (1993) Comparative protein modelling by satisfaction of spatial restraints. *J. Mol. Biol.*, 234, 779-815.
- Sali, A. (1995) Comparative protein modeling by satisfaction of spatial restraints. *Mol. Med. Today.* 1(6), 270-7.
- Sánchez, R., Sali, A. (1997) Evaluation of comparative protein structure modeling by MODELLER-3. *Proteins.* 1, 50-8.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) Molecular Cloning. *A Laboratory Manual.* Cold Spring Harbor Laboratory Press.
- Sanger, G.J., Nelson, D.R. (1989) Selective and functional 5-hydroxytryptamine 3 receptor antagonism by BRL 43694 (granisetron). *Europ. J. Pharmacol.* 159,113-124.

- Smart, O.S., Neduvilil, J.G., Wang, X., Wallace, B.A., Sansom, M.S. (1996) HOLE: a program for the analysis of the pore dimensions of ion channel structural models. *J. Mol. Graph.* 14(6), 354-60, 376.
- Sorensen, S.M., Humphreys, T.M. and Palfreyman, M.G. (1989) Effect of acute and chronic MDL 73,147EF, a 5-HT₃ receptor antagonist, on A9 and A10 dopamine neurons. *Europ. J. Pharm.* 163, 115-118.
- Spier, A.D., and Lummis, S.C.R. (2000) The Role of Tryptophan Residues in the 5-Hydroxytryptamine₃ Receptor Ligand Binding Domain. *J. Biol. Chem.*, 275, 5620–5625.
- Steward, L.J., Boess, F.G., Steele, J.A., Liu, D., Wong, N., and Martin, I.L. (2000) Importance of Phenylalanine 107 in Agonist Recognition by the 5-Hydroxytryptamine_{3A} Receptor. *Mol. Pharmacol.* 57, 1249-1255.
- Sullivan, N.L., Thompson, A.J., Price, K.L., Lummis, S.C.R. (2006) Defining the roles of Asn-128, Glu-129 and Phe-130 in loop A of the 5-HT₃ receptor. *Mol. Membr. Biol.*, 23, 442–451.
- Suryanarayanan, A., Joshi, P.R., Bikadi, Z., Mani, M., Kulkarni, T.R., Gaines, C., and Schulte, M.K. (2005) The loop C region of the murine 5-HT_{3A} receptor contributes to the differential actions of 5-hydroxytryptamine and m-chlorophenylbiguanide. *Biochem.* 44, 9140-9149.
- Tai, K., Fowler, P., Mokrab, Y., Stansfeld, P., and Sansom, S.P. (2008) Molecular Modeling and Simulation Studies of Ion Channel Structures, Dynamics and Mechanisms. *Meth. Cell Bio.* 90, 233-265.
- Tapia, L., Kuryatov, A., and Lindstrom, J. (2007) Ca²⁺ permeability of the (α₄)₂(β₂)₃ human acetylcholine receptors. *Mol. Pharmacol.* 71, 769-776.
- Tecott, L.H., Maricq, A.V. and Julius, D. (1993) Nervous system distribution of the serotonin 5-HT₃ receptor mRNA. *Proc. Natl. Acad. Sci. U S A.* 15, 90, 1430-4.
- Thompson, A.J., and Lummis, S.C.R. (2003). A single ring of charged amino acids at one end of the pore can control ion selectivity in the 5-HT₃ receptor. *British J. Pharm.*, 140, 359-365.
- Thompson, A.J., Price, K.L., Reeves, D.C., Chan, S.L., Chau P.L. and Lummis, S.C.R. (2005) Locating an Antagonist in the 5-HT₃ Receptor Binding Site Using Modeling and Radioligand Binding. *J. Biol. Chem.* 280, 20476–20482.
- Thompson, A.J., Chau, P.L., Chan, S.L., and Lummis, S.C.R. (2006) Unbinding pathways of an agonist and an antagonist from the 5-HT₃ Receptor. *Biophys. J.*, 90, 1979-1991.

- Thompson, A.J. and Lummis, S.C.R. (2006) 5-HT₃ Receptors. *Curr. Pharm. Des.* 12, 3615–3630.
- Thompson, A.J., Padgett, C.L., and Lummis, S.C. (2006) Mutagenesis and molecular modeling reveal the importance of the 5-HT₃ receptor F-loop. *J. Biol. Chem.* 281, 16576-16582.
- Thompson, A.J. and Lummis, S.C.R. (2007) The 5-HT₃ receptor as a therapeutic target. *Expert Opin. Ther. Targets* 2007, 11, 527–540.
- Thompson, A.J., Lochner, M., Lummis, S.C. (2008) Loop B is a major structural component of the 5-HT₃ receptor. *Biophys J.*, 95, 5728-36.
- Thompson, A.J., Lester, H.A., Lummis, S.C.R. (2010) The structural basis of function in Cys-loop receptors. *Biophysics*, 43(4), 449–499.
- Thompson, A. J., Price, K. L. and Lummis, S. C. R. (2011) Cysteine modification reveals which subunits form the ligand binding site in human heteromeric 5-HT₃AB receptors. *J. Physiol.* 589, 4243-57.
- Thompson, J.D., Higgins, D.G., Gibson, T.J., and Clustal, W. (1994). Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties, and weight matrix choice. *Nucleic Acids Res.*, 22, 4673-4680.
- Tsigenly, I., Sugiyama, N., Sine, S.M. and Taylor, P. (1997) A model of the nicotinic receptor extracellular domain based on sequence identity and residue location. *Biophys J.* 1997, 73, 52-56.
- Unwin, N. (2005) Refined structure of the nicotinic acetylcholine receptor at 4 Å resolution, *J. Mol. Biol.* 346, 967-989.
- Venkataraman, P., Joshi, P., Venkatachalan, S.P., Muthalagi, M., Parihar, H.S., Kirschbaum, K.S., Schulte, M.K. (2002) Functional group interactions of a 5-HT₃R antagonist *BMC Biochem.*, 3, 16.
- Vernekar, S.K.V., Hallaq, H.Y., Clarkson, G., Thompson, A.J., Silvestri, L., Lummis, S.C.R., and Lochner, M. (2010) Toward biophysical probes for the 5-HT₃ receptor: Structure-Activity Relationship Study of granisetron derivatives. *J. Med. Chem.*, 53, 2324-2328.
- Vriend, G. (1990). WHAT IF: A molecular modeling and drug design program. *J. Mol. Graph.*, 8, 52-56.

- Walstab, J., Rappold, G. and Niesler, B. (2010) 5-HT₃ receptors: Role in disease and target of drugs. *Pharmacol. Therap.* 128, 146-169.
- Wong, E.H.F., Clark, R., Leung, E., Loury, D., Bonhaus, D.W., Jakeman, L., Parnes, H., Whiting, R.L., Eglon, R.M. (1995) The interaction of RS 25259-197, a potent and selective antagonist, with 5-HT₃ receptors, in vitro. *British J. Pharmacol.* 114, 851-859.
- Yaakob, N., Malone, D.T., Exintaris, B., and Irving, H.R. (2011) Heterogeneity amongst 5-HT₃ receptor subunits: is this significant? *Curr. Mol. Med.*, 11, 57-68.
- Yakel, J.L., Lagrutta, A., Adelman, J.P., and North, R.A. (1993). Single amino acid substitution affects desensitization of the 5-hydroxytryptamine type 3 receptor expressed in *Xenopus* oocytes. *PNAS, USA* 90, 5030-5033.
- Yan, D., Schulte, M.K., Bloom, K.E., and White, M.M. (1999) Interaction of d-tubocurarine analogs with the 5-HT₃ receptor. *J. Biol. Chem.* 274, 5537-5541.
- Yan, D. and White, M.M. (2005) Spatial orientation of the antagonist granisetron in the ligand-binding site of the 5-HT₃ receptor. *Mol. Pharmacol.* 68, 365–371.
- Zhu, F., and Hummer, G. (2012) Theory and Simulation of Ion Conduction in the Pentameric GLIC Channel. *J. Chem. Theor. Comput.*, 8, 3759–3768.

List of my publications

- Cappelli, A., Manini, M., Paolino, M., Gallelli, A., Anzini, M., Mennuni, L., Del Cadia, M., De Rienzo, F., Menziani, M.C., and Vomero, S. (2011) Bivalent Ligands for the Serotonin 5-HT₃ Receptor. *ACS Med. Chem. Lett.*, 2 (8), 571–576.
- De Rienzo, F., Del Cadia, M., and Menziani, M.C. (2012) A first step towards the understanding of the 5-HT₃ receptor subunit heterogeneity from a computational point of view. *Phys. Chem. Chem. Phys.* 14, 12625–12636.
- Del Cadia, M., De Rienzo, F., Menziani, M.C. (2012) Approaching the 5-HT₃ Receptor heterogeneity by computational studies of the transmembrane and intracellular domains. *Submitted to Journal of Computer-Aided Molecular Design.*
- Del Cadia, M., De Rienzo, F., Weston, D.A., Thompson, J.A., Menziani, M.C., Lummis, S.C.R. (2012) Exploring a potential allosteric binding site of palonosetron in the 5-HT₃ receptor. *In preparation.*