

This is a pre print version of the following article:

Markers of Inflammation and Endothelial Dysfunction in Young Survivors from Acute Lymphoblastic Leukemia / Bruzzi, Patrizia; Bigi, Elena; Felici, Francesca; Lami, Francesca; Cano Garcinuno, Maria Del Carmen; Palazzi, Giovanni; Cellini, Monica; Predieri, Barbara; Iughetti, Lorenzo. - In: JOURNAL OF ADOLESCENT AND YOUNG ADULT ONCOLOGY. - ISSN 2156-5333. - 10:5(2021), pp. 599-605. [10.1089/jayao.2020.0114]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

01/05/2026 01:30

(Article begins on next page)

Journal of Adolescent and Young Adult Oncology

Journal Name: <http://mc.manuscriptcentral.com/jayao>

Markers of inflammation and endothelial dysfunction in young survivors from acute lymphoblastic leukemia.

Journal:	<i>Journal of Adolescent and Young Adult Oncology</i>
Manuscript ID	JAYAO-2020-0114.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Bruzzi, Patrizia; Azienda Ospedaliero-Universitaria di Modena Policlinico di Modena, Pediatric Unit</p> <p>Bigi, Elena; University of Modena and Reggio Emilia Faculty of Medicine and Surgery, Department of Medical and Surgical Sciences for Mothers, Children and Adults</p> <p>Felici, Francesca; University of Modena and Reggio Emilia Faculty of Medicine and Surgery, Post Graduate School of Pediatrics, Department of Medical and Surgical Sciences for Mothers, Children and Adults</p> <p>Lami, Francesca; Azienda Ospedaliero-Universitaria di Modena Policlinico di Modena, Pediatric Unit</p> <p>Cano Garcinuno, Maria del Carmen ; Azienda Ospedaliero-Universitaria di Modena Policlinico di Modena, Pediatric Oncology and Hematology Unit</p> <p>Palazzi, Giovanni; Azienda Ospedaliero-Universitaria di Modena Policlinico di Modena, Pediatric Oncology and Hematology Unit</p> <p>Cellini, Monica; Azienda Ospedaliero-Universitaria di Modena Policlinico di Modena, Pediatric Oncology and Hematology Unit</p> <p>Predieri, Barbara; University of Modena and Reggio Emilia Faculty of Medicine and Surgery, Department of Medical and Surgical Sciences for Mothers, Children and Adults</p> <p>Iughetti, Lorenzo; Università degli Studi di Modena e Reggio Emilia, Medical and Surgical sciences of Mother, Children and Adults</p>
Keyword:	Acute Lymphoblastic Leukemia, Biomarker, Chemotherapy, Late Effects, Pediatric, Survival
Manuscript Keywords (Search Terms):	cardiovascular disease, atherogenesis, inflammation, endothelial dysfunction, acute lymphoblastic leukemia, survivor
Abstract:	<p>Purpose: To assess subclinical markers of endothelial inflammation in young survivors from acute lymphoblastic leukemia (ALL) treated with chemotherapy without cranial irradiation.</p> <p>Methods: Anthropometric parameters [height (H), body mass index (BMI), waist circumference (WC), hip circumference (HC), WC/H and WC/HC ratio], blood pressure, lipid profile, serum markers of inflammation and endothelial dysfunction [Interleukin 6 (IL-6), Vascular Cell Adhesion Molecule (VCAM), Intercellular Adhesion Molecule (ICAM), Tumor Necrosis Factor-alfa (TNF-α), Endogenous secretory Receptor for Advanced Glycation Endproducts (Es-RAGE)] and carotid intima-media thickness (c-IMT) were assessed in a group of young ALL survivors and</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>in matched controls.</p> <p>Results: 28 ALL survivors (71% male, 18% prepubertal, aged 15.98±4.41 years, mean follow-up 8.57±3.14 years) exhibited lower levels of Es-RAGE than controls (0.18±0.07 vs. 0.27±0.08 ng/ml, p<0.001). Among survivors, Es-RAGE values significantly correlated with BMI-SD off-therapy (R2 -0.42), WC/H ratio (R2 -0.41), WC/HC ratio (R2 -0.38) and with low-density-lipoprotein cholesterol (LDL-C; R2 -0.43). Most of ALL survivors (78%) presented c-IMT above the 95th centile if compared with gender- and age-standard. Mean c-IMT value correlated with blood pressure (R2 0.56) and with LDL-C levels (R2 0.56). Metabolic syndrome (MetS) was fully detected only in one ALL survivor. Nevertheless, 18% ALL survivors presented more than one MetS diagnostic criteria: 14% insulin-resistance, 25% dyslipidemia and 17.8% hypertension.</p> <p>Conclusions: We demonstrated an initial functional vascular alteration in young ALL survivors even when treated with standard risk protocols. Our data already supports the activation at endothelial level of glycosylation and oxidation processes persistent long after the end of the treatment.</p>

SCHOLARONE™
Manuscripts

EXACT TITLE:

Markers of inflammation and endothelial dysfunction in young survivors from acute lymphoblastic leukemia.

AUTHORS

Patrizia Bruzzi, MD

Pediatric Unit, Department of Medical and Surgical Sciences for Mothers, Children and Adults, University of Modena and Reggio Emilia, Modena, 41124, Italy;

mail address: bruzzi.patrizia@aou.mo.it

Elena Bigi, MD

Pediatric Oncology and Hematology Unit, Department of Medical and Surgical Sciences for Mothers, Children and Adults, University of Modena and Reggio Emilia, Modena, 41124, Italy;

mail address: elena.bigi@gmail.com

Francesca Felici, MD

Post Graduate School of Pediatrics, Department of Medical and Surgical Sciences for Mothers, Children and Adults, University of Modena and Reggio Emilia, Modena, Italy

mail address: felici.francesca@ausl.re.it

Francesca Lami, MD

1
2
3 Pediatric Unit, Department of Medical and Surgical Sciences for Mothers, Children and Adults,
4 University of Modena and Reggio Emilia, Modena, 41124, Italy;

5
6
7
8 mail address: lami.francesca@aou.mo.it
9

10
11
12 Maria del Carmen Cano Garcinuno, MD
13

14 Pediatric Oncology and Hematology Unit, Department of Medical and Surgical Sciences for
15 Mothers, Children and Adults, University of Modena and Reggio Emilia, Modena, 41124, Italy;

16
17
18
19 mail address: cano.carmen@aou.mo.it
20
21
22

23
24 Palazzi Giovanni, MD
25

26 Pediatric Oncology and Hematology Unit, Department of Medical and Surgical Sciences for
27 Mothers, Children and Adults, University of Modena and Reggio Emilia, Modena, 41124, Italy;

28
29
30
31 mail address: palazzi.giovanni@aou.mo.it
32
33
34

35
36 Monica Cellini, MD
37

38 Pediatric Oncology and Hematology Unit, Department of Medical and Surgical Sciences for
39 Mothers, Children and Adults, University of Modena and Reggio Emilia, Modena, 41124, Italy;

40
41
42
43 mail address: cellini.monica@aou.mo.it
44
45
46

47
48 Barbara Predieri, MD, PhD

49 Pediatric Unit, Department of Medical and Surgical Sciences for Mothers, Children and Adults,
50 University of Modena and Reggio Emilia, Modena, 41124, Italy;

51
52
53
54 mail address: barbara.predieri@unimore.it
55
56
57
58
59
60

1
2
3
4
5
6 Lorenzo Iughetti, MD, PhD

7
8 Oncology and Hematology Pediatric Unit, Pediatric Unit, Post Graduate School of Pediatrics,
9
10 Department of Medical and Surgical Sciences for Mothers, Children and Adults, University of
11
12 Modena and Reggio Emilia, Modena, 41124, Italy;
13
14 mail address: lorenzo.iughetti@unimore.it
15
16
17
18

19 **Corresponding author:**

20
21 Patrizia Bruzzi, MD

22
23 Pediatric Unit, Department of Medical and Surgical Sciences for Mothers, Children and Adults,
24
25 University of Modena and Reggio Emilia, Modena, via Del Pozzo, n.41, 41124, Italy;
26
27

28 Phone number: +39-594224719

29
30 Fax number: +39-594223074

31
32
33 mail address: bruzzi.patrizia@aou.mo.it
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT:

Purpose: To assess subclinical markers of endothelial inflammation in young survivors from acute lymphoblastic leukemia (ALL) treated with chemotherapy without cranial irradiation.

Methods: Anthropometric parameters [height (H), body mass index (BMI), waist circumference (WC), hip circumference (HC), WC/H and WC/HC ratio], blood pressure, lipid profile, serum markers of inflammation and endothelial dysfunction [Interleukin 6 (IL-6), Vascular Cell Adhesion Molecule (VCAM), Intercellular Adhesion Molecule (ICAM), Tumor Necrosis Factor- α (TNF- α), Endogenous secretory Receptor for Advanced Glycation Endproducts (Es-RAGE)] and carotid intima-media thickness (c-IMT) were assessed in a group of young ALL survivors and in matched controls.

Results: 28 ALL survivors (71% male, 18% prepubertal, aged 15.98 ± 4.41 years, mean follow-up 8.57 ± 3.14 years) exhibited lower levels of Es-RAGE than controls (0.18 ± 0.07 vs. 0.27 ± 0.08 ng/ml, $p < 0.001$). Among survivors, Es-RAGE values significantly correlated with BMI-SD off-therapy ($R^2 -0.42$), WC/H ratio ($R^2 -0.41$), WC/HC ratio ($R^2 -0.38$) and with low-density-lipoprotein cholesterol (LDL-C; $R^2 -0.43$). Most of ALL survivors (78%) presented c-IMT above the 95th centile if compared with gender- and age-standard. Mean c-IMT value correlated with blood pressure ($R^2 0.56$) and with LDL-C levels ($R^2 0.56$). Metabolic syndrome (MetS) was fully detected only in one ALL survivor. Nevertheless, 18% ALL survivors presented more than one MetS diagnostic criteria: 14% insulin-resistance, 25% dyslipidemia and 17.8% hypertension.

Conclusions: We demonstrated an initial functional vascular alteration in young ALL survivors even when treated with standard risk protocols. Our data already supports the activation at endothelial level of glycosylation and oxidation processes persistent long after the end of the treatment.

BACKGROUND

Acute lymphoblastic leukemia (ALL) is the most common pediatric cancer with a 5-year survival rate of 90% in high-income countries.¹ Nowadays, therapeutic progresses in childhood and adolescence bring to high vulnerability later on in life and survival is still tempered by a wide variety of sequelae. An increase in the prevalence of metabolic syndrome (MetS) and of cardiovascular risk (CVR) over time is already well documented in survivors treated according to high-risk protocols including hematopoietic stem-cell transplantation (HSCT) conditioned with total-body irradiation (TBI) and/or cranial and/or abdominal radiotherapy (RT), regardless of body mass index (BMI).² Nevertheless, no homogeneous data are yet available on cardiovascular health and its risk of worsening in young ALL survivors treated with standard risk protocols without RT.³⁻⁴ Moreover, little is known about the molecular mechanisms leading to endothelial dysfunction and increase of CVR in this group of patients: cardiotoxicity could directly derived from the exposure to anthracyclines and local RT and, in addition, other endocrine dysfunctions could contribute to the development of MetS.

Several studies have already shown that early atherosclerotic lesions develop from morphological damages of the endothelium.⁵ When the normal physiology of the endothelium is altered, its permeability rises, its anti-thrombotic capacity reduces, the adhesion of leukocyte increases due to high levels of Vascular Cell Adhesion Molecule (VCAM) and Intercellular Adhesion Molecule (ICAM) and, at the same time, its tone augmented due to the reduced production of nitric oxide and the proliferation of vascular muscle cells.⁶ Interleukin 6 (IL-6) and Tumor Necrosis Factor- α (TNF- α) are primary inflammatory cytokines. IL-6 stimulates the production of acute phase proteins, amplifying the mechanisms of innate immunity and tissue remodelling.⁷ TNF- α influences each phase of the atherogenetic process: in fact, it promotes the

1
2
3 recruitment of monocytes inducing their adhesion to the endothelium and it could contribute to
4 the transformation and activation of macrophages.⁸ Advanced glycation end products (AGEs) are
5 complex and heterogeneous molecules derived *in vivo* from glycosylation and oxidation
6 processes. In addition to the membrane receptor, there is a soluble receptor, which is formed
7 either as a result of proteolytic cleavage of the membrane receptor (sRAGE) or by alternative
8 splicing of the mRNA that codes for RAGES (EsRAGE or endogenous secretory RAGE). In
9 particular conditions such as hyperglycemia, oxidative stress and inflammation, an increase of
10 AGEs and a concomitant reduction of EsRAGE have already been demonstrated.⁹ Previously,
11 only few studies measured these endothelium-derived blood vasoactive factors in similar cohorts
12 of childhood ALL survivors.¹⁰⁻¹¹

13
14
15
16
17
18
19
20
21
22
23
24
25
26 The evaluation of carotid intima-media thickness (c-IMT) is nowadays recommended by the
27 American Heart Association (AHA) as a non-invasive ultrasound parameter that identifies the
28 early stages of intimal wall alterations and, therefore, it represents an excellent indicator of the
29 pre-clinical stage of atherosclerosis also in childhood and adolescence.¹²

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
In the present study, we assess these subclinical markers of inflammation and endothelial
dysfunction in a uniform cohort of young survivors from ALL who underwent standard risk
therapeutic protocols without RT.

METHOD AND MATERIALS

Patients

Twenty-eight adolescents and young adults (93% Caucasian and 7% from South America)
survivors from childhood ALL treated locally (Oncology and Hematology Pediatric Unit,
Department of Medical and Surgical Sciences for Mothers, Children and Adults, University of
Modena and Reggio Emilia, Modena, Italy) according to ALL AIEOP 2000 and ALL AIEOP

2009 standard risk protocols¹³ (see Appendix A) and successfully off-therapy from at least 5 years were enrolled into the study (GROUP ALL, group A). Exclusion criteria counted the exposure to RT, HSCT and/or TBI, the presence of pre-existent conditions increasing CVR (eg. previous CV diseases, type 1 diabetes, familial hypercholesterolemia, genetic syndromes, smoking habit, family history of precocious cardiovascular disease in parents and grandparents...), the presence of one or more endocrine deficiencies and the use of other medications with potential effects on endothelial and CV functions (eg. growth hormone, inhaled corticosteroids, cholesterol lowering medications...).

Additionally, twenty-two age- and gender-matched healthy adolescents who met the same exclusion criteria worked as a control group (GROUP CONTROLS, group B). A written informed consent was obtained from all of them.

The study was approved by the Local Ethics Committee (code 56/13, approved on 24th June 2013).

Study design

This is an observational, transversal and case-control study.

The collection of anamnestic/therapeutic data together with an accurate anthropometric evaluation was performed in the group A during their follow up visit and in the group B.

Anthropometrical parameters [Height (H), body weight, body mass index (BMI), waist circumference (WC), hip circumference (HC), pubertal development] and blood pressure (BP) were evaluated according to standard procedures and expressed as mean \pm standard deviation (SD) with respect to chronological age and using national growth charts.¹⁴⁻¹⁷ Metabolic syndrome (MetS) comprises a cluster of CVR factors (hypertension, altered glucose metabolism,

dyslipidemia and abdominal obesity). In our cohort, MetS was defined according to Weiss' criteria.¹⁸

Blood samples were drawn early in the morning after overnight fasting. Moreover, group A underwent an evaluation of endothelial function through c-IMT assessment.

Laboratory methods

Plasma concentrations of total cholesterol (TC), low-density lipoprotein cholesterol (LDL-C), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C), fasting glucose and insulin were measured enzymatically using commercial kits (Cholesterol CHOD-PAD, HDL plus, Triglycerides GPO-PAP, Hitachi, Roche Diagnostic; Gluco-QuantR, Roche Diagnostic; Immulite 2000 system, Siemens Healthcare; respectively).

ICAM-1 and VCAM-1 levels were determined using OriGene company's ELISA kit (Human ICAM-1 and VCAM-1 ELISA kit). The lower limit of sensitivity was 10 pg/ml, with the reference range 156-10000 pg/ml. The levels of EsRAGE were determined through ELISA kit (B-Bridge International Company); normal values were 0.25-1.84 ng/ml. IL-6 and TNF- α concentrations were determined using ELISA kit (Diacclone Company). The lower limit of sensitivity was 2 pg/ml (reference range 6.25-200 pg/ml) and 10 pg/ml (reference range 25-800 pg/ml), respectively.

Assessment of c-IMT

A single experienced sonographer performed all the measurements after a fasting night, in a quiet and temperature-controlled room. An ultrasound system (ESAOTE, MyLab70XVision) equipped with vascular software for two-dimensional (2D) imaging, color Doppler and high-frequency vascular transducer (8-15 MHz) was used. This performance followed standardized protocols and results were interpreted according to pediatric standard.¹⁹

Statistical Analysis

All variables were expressed as mean \pm SD. The comparisons among clinical and biochemical data between groups were performed using *U* Mann-Whitney test (STATISTICA, StatSoft Inc., Tulsa, OK, USA). Univariate analysis was performed to correlate variables and Pearson's *r* coefficient was calculated. Statistical significance was determined at a *p* value of < 0.05 .

RESULTS

Clinical and metabolic features

Table 1 shows the main clinical characteristics of all enrolled subjects.

Groups A and B did not differ in age, anthropometrical parameters and rates of distribution of gender, pubertal status and ethnicity, as shown in Table 1.

In Group A, BMI SD significantly increased from diagnosis of ALL to study time (off-therapy). Nevertheless, fasting glycemetic metabolism [fasting glucose/insulin ratio 13.03 ± 6.71 (range 2.23 – 28.51); HOMA index 1.96 ± 1.54 (range 0.5 – 8.47)] and lipid parameters [TC 155.41 ± 24.84 mg/dl (range 99-193); LDL-C 90.28 ± 21.76 mg/dl (range 47-134); HDL-C 55.96 ± 14.21 (range 37 – 89); TG 70.77 ± 39.74 mg/dl (range 32-189)] were within the normal range.

According to Weiss' definition¹⁸, metabolic syndrome (MetS) was fully detected only in one ALL survivor. Nevertheless, 18% ALL survivors presented more than one MetS diagnostic criteria: 14% (4/28) showed insulin-resistance, 25% dyslipidemia (7/28), 17.8% hypertension (5/28) and 42% (12/28) abdominal obesity.

Inflammation and vascular parameters

Group A exhibited lower levels of Es-RAGE than group B (Table 2). No other differences in inflammation and vascular markers were detected between groups (Table 2). Among survivors, Es-RAGE values inversely correlated with BMI-SD off-therapy ($R^2 -0.42$), WC/H ratio ($R^2 -$

0.41), WC/HC ratio (R^2 -0.38) and with LDL-C values (R^2 -0.43) (Table 3). IL-6 levels directly correlated with WC/H ratio (R^2 0.41), WC/HC ratio (R^2 0.51) and TG values (R^2 0.40) while TNF- α values correlated with diastolic blood pressure (DBP; R^2 0.50) (Table 3).

Ultrasound parameter – c IMT

In group A, mean c-IMT is located above the 95th age-standard centile (mean value 0.55 ± 0.14 mm, range 0.4-0.85).¹⁹ Even when single data were analyzed, in 70% of males and in all females, c-IMT was above the 95th centile if compared to gender- and age-standard.¹⁹ In group A, mean c-IMT correlated with both systolic (R^2 0.56) and diastolic blood pressure (R^2 0.66) and LDL-C levels (R^2 0.56)(Table 3).

DISCUSSION

Clinical and metabolic features

Treatment-related risk factors for obesity and MetS in ALL survivors have been previously reviewed in literature²⁰ and extensively by our group.²¹ In 2014, we have already demonstrated that, among 107 Caucasian ALL patients treated on AIEOP protocols without radiotherapy between 1989 and 2000, BMI-SD increased persistently from the start of chemotherapy until the achievement of final height.²² As current data shown, we confirmed the increase of BMI-SD from ALL diagnosis through a long-term follow-up. Despite BMI-SD trend, in group A, at study time, mean values of BMI-SD, WC/H and WH/HC ratios, glucose and lipid profile were within the normal age-standard (table 1). Nevertheless, our population is not considerable as “metabolic healthy”: the 3.5% (1/28) showed MetS, 14% (4/28) insulin-resistance, 25% dyslipidemia (7/28), 17.8% hypertension (5/28) and 42% (12/28) abdominal obesity. Similar findings are presented in other studies of adult survivors of childhood ALL even if prevalence rates for components of

1
2
3 MetS vary widely in literature, due to heterogeneity in diagnostic criteria and study reporting
4
5 methods.²³⁻²⁴
6

7 8 **Inflammation and vascular parameters** 9

10 Our data show that, after a long follow-up period since the end of standard chemotherapy, even
11 without previous exposition to high-risk therapies (as TBI and/or HSCT), childhood ALL
12 survivors showed a pathological c-IMT and a marked increase in Es-RAGE consumption in
13 comparison to age-matched healthy controls. In line with our results, Felicetti F et al. recently
14 reported increased plasmatic levels of AGEs in a cohort of 18 adult survivors of childhood ALL
15 treated with HSCT conditioned with TBI.² As already known, AGEs may contribute to the
16 beginning and the development of vascular lesions, attracting monocytes to bind to the vessel
17 surface, followed by their migration into the vessel wall and subsequent activation of endothelial
18 cells and production of other mediators. In Felicetti's study, survivors showed also higher levels
19 of C-reactive protein (CRP), IL-1 α , and IL-17 than controls, suggesting the presence of chronic
20 inflammatory state, in agreement with other study.²⁵ In our study only esRAGE but no other
21 inflammatory molecules differed between ALL survivors and controls. Considering the central
22 role of AGEs in glycoxidation of lipoproteins and foam cells formation, we can speculate that
23 glycosylation and oxidation processes are more expressed in this category of patients at that age
24 (adolescence) than inflammation itself. The latter could be more pronounce later on in life
25 (adulthood). An unbalanced oxidative stress could be not only the trigger but also a persistent
26 and silent disorder leading to inflammation and endothelium dysfunction: this is the so called
27 "inflammaging" process already described in cancer survivors.²⁶ In addition to a well-known
28 radiation-induced damage, also chemotherapy agents alone, such as anthracyclines, cause an
29 immediate increase in the production of reactive oxygen species (ROS), generating a persistent
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

oxidative stress²⁷ and a direct apoptosis of vascular endothelial cells²⁸ and high dose of corticosteroids could increase visceral adiposity and reduce insulin-sensitivity.¹⁰ Our patients were exposed to a cumulative mean dose of anthracyclines of 244 ± 3.14 mg/m² and prednisone of 926 ± 42 mg/m² (*data not shown*). In the study performed by Sadurska et al in 2018 among an heterogeneous group of 64 childhood ALL survivors (70.3% treated with only chemotherapy, 70% exposed to a cumulative dose of anthracyclines less than 240 mg/m²), blood concentration of sICAM, IL-6 and high-sensitivity C-reactive protein as well as cIMT values were higher in ALL group than controls.¹¹ We can only speculate that dissimilarities between our and these results derived from differences in chemotherapeutic protocols and from the exposition of about 30% of survivors to high-risk therapy, such as RT, in Sadurska's cohort. Nevertheless, endothelial damage appears to be persistent.

In summary, ALL survivors, even if treated only with chemotherapy, have to be considered as unhealthy metabolic subjects similar to their obese and overweight peers.

Ultrasound parameter – c IMT

Among ALL survivors, we demonstrated a morphological alteration of the vascular wall.

Our results seem far from a recent study performed among 81 childhood ALL survivors aged 5-25 years (mean follow-up length 5.7 ± 3.9 years).²⁹ In this cohort, mean cIMT value was not statistically different between childhood ALL survivors and controls (0.43 ± 0.03 vs 0.42 ± 0.03 , respectively).²⁹ Discrepancies could be justified by the age of survivors (older in our study) and by the length of follow-up after the end of chemotherapy (longer in our study). Moreover, in this recent publication, no details about ALL protocols were provided and thus, any speculation in comparing our and their results seems to be inconsistent.²⁹ In 2016, Giordano et al demonstrated that childhood ALL survivors had lower FMD of the brachial artery but not pathological cIMT

1
2
3 when compared to healthy controls.⁴ As reported in literature in other categories of patients with
4 an high CV risk as children and adolescents with heterozygous familial hypercholesterolemia,
5 alterations in FMD are considered earlier indicators of endothelial dysfunction than cIMT.³⁰ In
6 line with this finding, when our and Giordano's data are compared, our population presented as
7 older (15.9±4.4 vs. 9.5±4.1 years old, respectively) and the follow-up as longer (average 8.5±3.1
8 vs. 1.17 years, respectively) than Giordano's ones.⁴ Moreover, their study population comprises
9 52 childhood ALL survivors treated with different risk protocols (6 high-risk, 29 intermediate
10 risk, 17 standard risk; 8 patients underwent cranial RT).⁴ The importance of the timing (age)
11 when cIMT is performed, is confirmed by Sadurska and colleagues: in their study, ALL
12 survivors, similar to our cohort in terms of age (15.5±5.5 years old) and length of follow-up (9.2
13 ± 4.4 years), presented an increased cIMT than controls.¹¹

14
15
16
17
18
19
20
21
22
23
24
25
26
27
28 In the already mentioned Ociepa's study, a significant correlation between 24-hour systolic BP
29 SD and cIMT-SD in ALL survivors was documented.²⁹ This seems to be in line with our results
30 demonstrating a direct correlation between mean c-IMT and both systolic and diastolic blood
31 pressure and LDL-C levels (Table 3). Other study are in line with our results.³¹ Moreover, in
32 long-term ALL survivors, endothelial dysfunction, measured by both FMD and cIMT, seems to
33 be reversible and counteracted with physical activity as a simple home-based exercise program³²⁻
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1
2
3 system leading them to 4 times greater risk of developing anthracycline-induced clinical heart
4 failure than male cancer survivors.³³ In our study, 70% of male and all female ALL survivors
5 presented a pathological c-IMT and we did not observe sex differences in BMI values, blood
6 pressure, lipid profile and serum markers of inflammation and endothelial dysfunction (*data not*
7 *shown*). Considering the exiguity of our population as the main study's limitation, the influence
8 of gender needs to be analyzed in larger cohorts.

16 CONCLUSION

19 Our study demonstrated that even standard risk childhood ALL survivors, treated on modern
20 protocols, presented precocious functional and structural endothelial alterations and we support
21 the use of cIMT in screening such complications at least in adolescence. **We hope to shortly**
22 **reinforce present data measuring cIMT also in healthy controls: this improvement is now**
23 **limited by the current Covid19 emergency.**

31 In our opinion, considering the state-of-art, the measurement of serum inflammation and vaso-
32 active molecules (e.g. Es-RAGE, IL-6, VCAM, ICAM and TNF- α) and their interpretation
33 seems more challenging in the clinical practice: their meanings and sensitivity could be
34 influenced by many biochemical factors that still have to be identified leading to a clarification
35 of CV etio-pathogenetic mechanisms.

42 Currently, our data support the inclusion in the long-term survivor systematic follow-up
43 programs of an early and routinely evaluation of both standard metabolic parameters (e.g. BMI,
44 WC/H ratio, WC/HC ratio, BP, glucose and lipid profile) and endothelial function through non-
45 invasive useful tools (FMD and c-IMT). This addition to current practice seems fundamental for
46 providing early detection and treatment of potentially serious late-onset CV complications,
47 which can be ameliorated by lifestyle modifications³⁴. Therefore, the counseling and promotion

1
2
3 of healthy lifestyles are important aspects in the long-term follow-up care and must be encourage
4 soon after the end of treatment.³⁵ Screening and prevention could, therefore, partially reverse the
5
6 course.
7
8
9

10 11 12 ACNOWLEDGMENTS

13
14 We thank all the study participants and their families for their availability and attitude.

15
16
17 **Preliminary data of our study have been presented as an abstract and a rapid free**
18
19 **communication at the 58th Annual European Society for Paediatric Endocrinology (ESPE)**
20
21 **Meeting organized in Wien (Austria Center), Austria, 19th - 21st September 2019.**
22

23 24 AUTHOR DISCLOSURE STATEMENT

25
26 No competing financial interests exist.

27 28 FUNDING INFORMATION

29
30 The authors received no specific funding for this work.

31 32 REFERENCES:

- 33
34
35 1. Smith MA, Seibel NL, Altekruse SF, et al. Outcomes for children and adolescents with cancer:
36 Challenges for the twenty- first century. J Clin Oncol 2010;28:2625–2634.
37
38 2. Felicetti F, Cento AS, Fornengo P, et al. Advanced glycation end products and chronic
39 inflammation in adult survivors of childhood leukemia treated with hematopoietic stem cell
40 transplantation. Pediatr Blood Cancer 2020;67(3):e28106.
41
42 3. Chemaitilly W, Cohen LE, Mostoufi-Moab S, et al. Endocrine Late Effects in Childhood
43 Cancer Survivors. J Clin Oncol 2018;36 (21):2153-2159.
44
45 4. Giordano P, Muggeo P, Delvecchio M, et al. Endothelial dysfunction and cardiovascular risk
46 factors in childhood acute lymphoblastic leukemia survivors. Int J Cardiol 2017;228: 621-627.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 5. Shaul PW. Role of the Endothelium in the Metabolic Syndrome: IIB or Not IIB. *Am J Med*
4
5 *Sci* 2015;349 (1):3–5.
6
- 7
8 6. Paulus P, Jennewein C, Zacharowski K. Biomarkers of endothelial dysfunction: can they help
9
10 us deciphering systemic inflammation and sepsis? *Biomarkers* 2011; 16 S1: S11-21.
11
- 12 7. Woods A, Brull DJ, Humphries SE, Montgomery HE. Genetics of inflammation and risk of
13
14 coronary artery disease: the central role of interleukin-6. *Eur Heart J* 2000; 21:1574-83.
15
- 16 8. Kleinbongard P, Heusch G, Schulz R. TNF α in atherosclerosis, myocardial
17
18 ischemia/reperfusion and heart failure. *Pharmacol Ther* 2010;295-314.
19
- 20 9. Basta G, Sironi AM, Lazzarini G, et al. Circulating soluble receptor for advanced glycation
21
22 end products is inversely associated with glycemic control and S100A12 protein. *J Clin*
23
24 *Endocrinol Metab* 2006; 91: 4628-34.
25
26
- 27 10. Sulicka J, Surdacki A, Mikolajczyk T, et al. Elevated markers of inflammation and
28
29 endothelial activation and increased counts of intermediate monocytes in adult survivors of
30
31 childhood acute lymphoblastic leukemia. *Immunobiology* 2013; 218 (5): 810-816.
32
33
- 34 11. Sadurska E, Zaucha-Prazmo A, Brodzisz A, et al. Premature atherosclerosis after treatment
35
36 for acute lymphoblastic leukemia in childhood. *Ann Agric Environ Med* 2018; 25(1): 71-76.
37
38
- 39 12. Dalla Pozza R, Ehringer-Schetitska D, Fritsch P, et al. Intima media thickness measurement
40
41 in children: a statement from the Association for European Paediatric Cardiology (AEPC)
42
43 working group on cardiovascular prevention endorsed by the association for European Paediatric
44
45 Cardiology. *Atherosclerosis* 2015; 238 (2): 380-387.
46
47
- 48 13. Testi AM, Attarbaschi A, Valsecchi MG, et al; AIEOP-BFM (Associazione Italiana di
49
50 *Ematologia e Oncologia Pediatrica & Berlin-Frankfurt-Münster*) Study Group. Outcome of
51
52 adolescent patients with acute lymphoblastic leukaemia aged 10-14 years as compared with those
53
54
55
56
57
58
59
60

aged 15-17 years: Long-term results of 1094 patients of the AIEOP-BFM ALL 2000 study. *Eur J Cancer* 2019;122: 61-71.

14. Cacciari E, Milani S, Balsamo A, et al. Italian cross-sectional growth charts for height, weight and BMI (2 to 20 yr). *J Endocrinol Invest* 2006; 29: 581–593.

15. Zannolli R, Morgese G. Waist percentiles: a simple test for atherogenic disease? *Acta Paediatr* 1996; 85: 1368–1369.

16. Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. *Arch Dis Child* 1969; 44: 291–303.

17. Baker-Smith CM, Flinn SK, Flynn JT, et al; SUBCOMMITTEE ON SCREENING AND MANAGEMENT OF HIGH BP IN CHILDREN. Diagnosis, Evaluation, and Management of High Blood Pressure in Children and Adolescents. *Pediatrics* 2018;142(3).

18. Weiss R, Bremer AA, Lustig RH. What is metabolic syndrome, and why are children getting it? *Ann N Y Acad Sci* 2013;1281:123-40.

19. Doyon A, Kracht D, Bayazit AK, et al. Carotid artery intima – media thickness and distensibility in children and adolescents: reference values and role of body dimensions. *Hypertension* 2013; 62:550 – 556.

20. Rosen GP, Nguyen HT, Shaibi GQ. Metabolic Syndrome in Pediatric Cancer Survivors: A Mechanistic Review. *Pediatr Blood Cancer* 2013; 60:1922–1928.

21. Iughetti L, Bruzzi P, Predieri B, Paolucci P. Obesity in patients with acute lymphoblastic leukemia in childhood. *Ital J Pediatr* 2012; 38:4.

22. Bruzzi P, Predieri B, Corrias A, et al. Final height and body mass index in adult survivors of childhood acute lymphoblastic leukemia treated without cranial radiotherapy: a retrospective longitudinal multicenter Italian study. *BMC Pediatr* 2014;14:236.

- 1
2
3 23. Gibson TM, Ehrhardt MJ, Ness KK. Obesity and Metabolic Syndrome Among Adult
4 Survivors of Childhood Leukemia. *Curr Treat Options Oncol* 2016; 17(4):17.
5
6
7 24. Levy E, Samoilenko M, Morel S, et al. Cardiometabolic Risk Factors in Childhood,
8 Adolescent and Young Adult Survivors of Acute Lymphoblastic Leukemia – A Petale Cohort.
9
10
11
12
13
14
15 25. Ariffin H, Azanan MS, Abd Ghafar SS, et al. Young adult survivors of childhood acute
16 lymphoblastic leukemia show evidence of chronic inflammation and cellular aging. *Cancer*
17
18
19
20
21
22 26. Aragno M, Mastrocola R. Dietary sugars and endogenous formation of advanced glycation
23 endproducts: emerging mechanisms of disease. *Nutrients* 2017; 9(4): 385.
24
25
26 27. Wu S, KO YS, Teng MS, et al. Adriamycin-induced cardiomyocyte and endothelial cell
27 apoptosis: in vitro and in vivo studies. *J Mol Cell Cardiol* 2002; 34(12): 1596-1607.
28
29
30
31 28. Dengel DR, Ness KK, Glasser SP, et al. Endothelial function in young adult survivors of
32 childhood acute lymphoblastic leukemia. *J Pediatr Hematol Oncol* 2008; 30(1): 20-25.
33
34
35 29. Ociepa T, Posio W, Sawicki M, Urasinski T. CIMT does not identify early vascular changes
36 in childhood acute lymphoblastic leukemia survivors. *Adv Clin Exp Med* 2020; 29 (2): 243-249.
37
38
39
40 30. Vlahos AP, Naka KK, Bechlioulis A, et al. Endothelial dysfunction, but not structural
41 atherosclerosis, is evident early in children with heterozygous familial hypercholesterolemia.
42
43
44
45
46
47 31. Dawson JD, Sonka M, Blecha MB, et al. Risk factors associated with aortic and carotid
48 intima-media thickness in adolescents and young adults: The Muscatine Offspring Study. *J Am*
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 32. Jarvela LS, Niinikoski H, Heinonen OJ, et al. Endothelial function in long-term survivors of
4 childhood acute lymphoblastic leukemia: effects of a home-based exercise program. *Pediatr*
5
6 *Blood Cancer* 2013; 60 (9):1546-1551.
7
8
9
10 33. Caru M, Curnier D. Sex and gender considerations after surviving acute lymphoblastic
11
12 leukemia: an exercise oncology context. *J Adolesc Young Adult Oncol* 2020; 9(3):441-444.
13
14 34. Oeffinger KC, Mertens AC, Sklar CA, et al. Chronic health conditions in adult survivors of
15
16 childhood cancer. *N Engl J Med* 2006; 355 (15):1572-82.
17
18
19 35. American Academy of Pediatrics Section on Hematology/Oncology Children's Oncology
20
21 Group. Long-term Follow-up Care for Pediatric Cancer Survivors. *Pediatrics* 2009; 123(3):906-
22
23 15.
24
25 36. Scott JM, Li N, Liu Q, et al. Association of exercise with mortality in adult survivors of
26
27 childhood cancer. *JAMA Oncol* 2018; 4: 1352-58.
28
29
30 37. Muggeo P, Muggeo VMR, Giordano P, et al. Cardiovascular dysfunction and vitamin D
31
32 status in childhood acute lymphoblastic leukemia survivors. *World J Pediatr* 2019; 15(5):465-
33
34 470.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES:

Table 1. Clinical features of total study population (Group A and B).

Data are presented as mean value \pm SD (range)

Legend: ALL, Acute Lymphoblastic Leukemia; BMI, Body Mass Index; N.A., Not Applicable; SD, Standard Deviation; TH, Target Height; WC/H, Waist circumference/Height; WC/HC, Waist circumference/Hip circumference.

* $p < 0.05$ at diagnosis ALL vs. off-therapy

Parameters	Group A (n.28)	Group B (n.22)	p-value
Age (years)	15.98 \pm 4.41 (9.18 – 24.88)	16.59 \pm 5.60 (6.00 – 27.13)	0.48
Male (%)	71	64	0.45
Prepubertal (%)	18	25	0.35
Caucasian (%)	93	92	0.60
Follow-up (years)	8.57 \pm 3.14 (4.05 – 15.69)	N.A.	N.A.
Height SD at diagnosis ALL	0.15 \pm 1.27 (-2.66 – 2.86)	N.A.	N.A.
Height SD off-therapy	0.01 \pm 1.05 (-1.82 – 2.19)	0.09 \pm 0.98 (-1.45 – 1.88)	0.09
Height SD off-therapy TH-adjusted	0.26 \pm 0.84 (-1.80 – 1.65)	N.A.	N.A.
BMI SD at diagnosis ALL	-0.34 \pm 1.42 (-3.35 – 2.80)	N.A.	N.A.
BMI SD off-therapy	0.55 \pm 1.30 (-1.25 – 3.13)*	0.42 \pm 1.01 (-1.33 – 2.34)	0.37
WC/H ratio off-therapy	0.48 \pm 0.08 (0.39 – 0.71)	N.A.	N.A.
WC/HC ratio off-therapy	0.91 \pm 0.07 (0.73 – 1.06)	N.A.	N.A.

Table 2. Serum endothelial biomarkers in both groups.

Data are presented as mean value \pm SD.

Legend: Es-RAGE, Endogenous secretory Receptor for Advanced Glycation Endproducts; ICAM, Intercellular Adhesion Molecule; IL-6, Interleukin 6; SD, Standard Deviation; TNF- α , Tumor Necrosis Factor-alfa; VCAM, Vascular Cell Adhesion Molecule.

Parameters	Group A	Group B	p- value
IL-6 (pg/ml)	1.39 \pm 0.74	1.70 \pm 1.24	0.67
VCAM (pg/ml)	1488.92 \pm 534.53	1614.31 \pm 725.11	0.65
ICAM (pg/ml)	109.96 \pm 55.3	104.30 \pm 55.63	0.46
TNF- α (pg/ml)	8.15 \pm 7.39	7.78 \pm 5.15	0.75
Es-RAGE (ng/ml)	0.18 \pm 0.07	0.27 \pm 0.08	< 0.001

Table 3. Correlation analysis (R^2) between cardiovascular risk markers and clinical and biochemical parameters in Group A (* $p < 0.05$).

Legend: BMI, Body Mass Index; c-IMT, carotid-Intima Media Thickness; DBP, Diastolic Blood Pressure; Es-RAGE, Endogenous secretory Receptor for Advanced Glycation Endproducts; IL-6, Interleukin 6; LDL-C, Low-density Lipoprotein- Cholesterol; SBP, Systolic Blood Pressure; SD, Standard Deviation; TG, Triglycerides; TNF- α , Tumor Necrosis Factor-alfa; WC/H, Waist circumference/Height; WC/HC, Waist circumference/Hip circumference.

Parameters	Es-Rage	IL-6	TNF- α	c-IMT
BMI-SDS off-therapy	-0.42*	+0.32	+0.02	+0.40
WC/H ratio	-0.41*	+0.41*	+0.12	+0.36
WC/HC ratio	-0.38*	+0.51*	+0.20	-0.03
LDL-C (mg/dl)	-0.43*	+0.08	-0.21	+0.56*
TG (mg/dl)	-0.33	+0.40*	+0.04	+0.39
SBP (mmHg)	-0.16	-0.02	+0.04	+0.56*
DBP (mmHg)	+0.13	+0.11	+0.50*	+0.66*

Appendix A.

AIEOP ALL 2000 and 2009 standard risk protocol (cumulative doses expressed in mg/m²).

	Chemotherapy: Phases			
	Patients Group A N (%)	Induction	Consolidation	Reinduction
AIEOP ALL 2000	23 (82%)	63 days PDN (1680 + tapering) VCR (6) DNR (120) L-ASP (40000) CPM (2000) ARA-C (1200) 6MP (1680)	56 days MTX (8000) 6MP (1400)	49 days DXM (210 + tapering) VCR (6) ADM (120) L-ASP (40000) CPM (1000) ARA-C (600) 6TG (840)
AIEOP ALL 2009	5 (18%)	63 days PDN (1680 + tapering) VCR (6) DNR (120) PEG-ASP (5000) CPM (2000) ARA-C (1200) 6MP (1680)	56 days MTX (20000) 6MP (1400)	49 days DXM (210 + tapering) VCR (6) ADM (120) PEG-ASP (2500) CPM (1000) ARA-C (600) 6TG (840)

Legend: PDN, Prednisone; DXM, dexamethasone; VCR, vincristine; DNR, daunorubicin; ADM, Adriamycin; CPM, cyclophosphamide; ARA-C, cytarabine; L-ASP, l-asparaginase (cumulative dose expressed in UI/m²); PEG-ASP, pegylated asparaginase; 6MP, mercaptopurine; 6TG, thioguanine; MTX, methotrexate

Table 1. Clinical features of total study population (Group A and B).

1 Data are presented as mean value ± SD (range)
2
3

4 **Legend:** ALL, Acute Lymphoblastic Leukemia; BMI, Body Mass Index; N.A., Not Applicable;
5 SD, Standard Deviation; TH, Target Height; WC/H, Waist circumference/Height; WC/HC,
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41

* p<0.05 at diagnosis ALL vs. off-therapy

Parameters	Group A (n.28)	Group B (n.22)	p-value
Age (years)	15.98 ± 4.41 (9.18 – 24.88)	16.59 ± 5.60 (6.00 – 27.13)	0.48
Male (%)	71	64	0.45
Prepubertal (%)	18	25	0.35
Caucasian (%)	93	92	0.60
Follow-up (years)	8.57 ± 3.14 (4.05 – 15.69)	N.A.	N.A.
Height SD at diagnosis ALL	0.15 ± 1.27 (-2.66 – 2.86)	N.A.	N.A.
Height SD off-therapy	0.01 ± 1.05 (-1.82 – 2.19)	0.09 ± 0.98 (-1.45 – 1.88)	0.09
Height SD off-therapy TH-adjusted	0.26 ± 0.84 (-1.80 – 1.65)	N.A.	N.A.
BMI SD at diagnosis ALL	-0.34 ± 1.42 (-3.35 - 2.80)	N.A.	N.A.
BMI SD off-therapy	0.55 ± 1.30 (-1.25 - 3.13)*	0.42 ± 1.01 (-1.33 – 2.34)	0.37
WC/H ratio off-therapy	0.48 ± 0.08 (0.39 – 0.71)	N.A.	N.A.
WC/HC ratio off-therapy	0.91 ± 0.07 (0.73 – 1.06)	N.A.	N.A.

Table 2. Serum endothelial biomarkers in both groups.

Data are presented as mean value \pm SD.

Legend: Es-RAGE, Endogenous secretory Receptor for Advanced Glycation Endproducts; ICAM, Intercellular Adhesion Molecule; IL-6, Interleukin 6; TNF- α , Tumor Necrosis Factor- α ; SD, Standard Deviation; VCAM, Vascular Cell Adhesion Molecule.

Parameters	Group A	Group B	p- value
IL-6 (pg/ml)	1.39 \pm 0.74	1.70 \pm 1.24	0.67
VCAM (pg/ml)	1488.92 \pm 534.53	1614.31 \pm 725.11	0.65
ICAM (pg/ml)	109.96 \pm 55.3	104.30 \pm 55.63	0.46
TNF- α (pg/ml)	8.15 \pm 7.39	7.78 \pm 5.15	0.75
Es-RAGE (ng/ml)	0.18 \pm 0.07	0.27 \pm 0.08	< 0.001

Table 3. Correlation analysis (R^2) between cardiovascular risk markers and clinical and biochemical parameters in Group A (* $p < 0.05$).

Legend: BMI, Body Mass Index; c-IMT, carotid-Intima Media Thickness; DBP, Diastolic Blood Pressure; Es-RAGE, Endogenous secretory Receptor for Advanced Glycation Endproducts; IL-6, Interleukin 6; LDL-C, Low-density Lipoprotein- Cholesterol; SBP, Systolic Blood Pressure; SDS, Standard Deviation Score; TG; Triglycerides; TNF- α , Tumor Necrosis Factor-alfa; WC/H, Waist circumference/Height; WC/HC, Waist circumference/Hip circumference.

Parameters	Es-Rage	IL-6	TNF- α	c-IMT
BMI-SDS off-therapy	-0.42*	+0.32	+0.02	+0.40
WC/H ratio	-0.41*	+0.41*	+0.12	+0.36
WC/HC ratio	-0.38*	+0.51*	+0.20	-0.03
LDL-C (mg/dl)	-0.43*	+0.08	-0.21	+0.56*
TG (mg/dl)	-0.33	+0.40*	+0.04	+0.39
SBP (mmHg)	-0.16	-0.02	+0.04	+0.56*
DBP (mmHg)	+0.13	+0.11	+0.50*	+0.66*

Appendix A.**AIEOP ALL 2000 and 2009 standard risk protocol (cumulative doses expressed in mg/m²).**

	Patients Group A N (%)	Chemotherapy: Phases		
		Induction	Consolidation	Reinduction
AIEOP ALL 2000	23 (82%)	63 days PDN (1680 + tapering) VCR (6) DNR (120) L-ASP (40000) CPM (2000) ARA-C (1200) 6MP (1680)	56 days MTX (8000) 6MP (1400)	49 days DXM (210 + tapering) VCR (6) ADM (120) L-ASP (40000) CPM (1000) ARA-C (600) 6TG (840)
AIEOP ALL 2009	5 (18%)	63 days PDN (1680 + tapering) VCR (6) DNR (120) PEG-ASP (5000) CPM (2000) ARA-C (1200) 6MP (1680)	56 days MTX (20000) 6MP (1400)	49 days DXM (210 + tapering) VCR (6) ADM (120) PEG-ASP (2500) CPM (1000) ARA-C (600) 6TG (840)

Legend: PDN, Prednisone; DXM, dexamethasone; VCR, vincristine; DNR, daunorubicin; ADM, Adriamycin; CPM, cyclophosphamide; ARA-C, cytarabine; L-ASP, l-asparaginase (cumulative dose expressed in UI/m²); PEG-ASP, pegylated asparaginase; 6MP, mercaptopurine; 6TG, thioguanine; MTX, methotrexate