

This is the peer reviewed version of the following article:

Proposed roles of the immune response regulator-ThPOK in human colorectal cancer progression / Sena, Paola; Mariani, Francesco; Roncucci, Luca; Benincasa, Marta; Ponz De Leon, Maurizio; Palumbo, Carla. - In: ITALIAN JOURNAL OF ANATOMY AND EMBRYOLOGY. - ISSN 1122-6714. - STAMPA. - 117 - n. 2 (suppl.):(2012), pp. 177-177. (66h Meeting of the Italian Society of Anatomy and Histology Pistoia (Italy) 20-23 September, 2012).

Firenze University Press
Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

03/05/2026 08:26

(Article begins on next page)

Proposed roles of the immune response regulator- ThPOK in human colorectal cancer progression

Paola Sena¹, Francesco Mariani², Luca Roncucci², Marta Benincasa¹, Maurizio Ponz de Leon² and Carla Palumbo¹

¹ Dipartimento di Scienze Biomediche, Metaboliche e Neuroscienze, Sezione di Morfologia Umana, Università di Modena e Reggio Emilia, Modena, Italy

² Dipartimento di Medicina e Specialità Mediche, Policlinico Universitario, Modena, Italy

Solid tumours are commonly infiltrated by several immune cells [1-3]. In cancer, immune cells play conflicting roles with both the potentials to eliminate or to promote malignancy. In contrast to infiltration of cells responsible for chronic inflammation, the presence of high numbers of lymphocytes, especially T cells, has been reported to be important as indicator of good prognosis in many types of cancer [4-7]. The thorough knowledge of both manners and pathways with which tumors are able to evade immune-mediated attack, once established, is therefore of crucial importance. The strategies to escape anti-tumor immune responses include the limited priming or differentiation of antitumor T cells and the role of tumor microenvironment in order to prevent infiltration or activation of effector phase functions. We proposed to evaluate the role of Th inducing POZ-Kruppel Factor (ThPOK), a transcriptional regulator of T cell fate, in tumour-induced immune system plasticity during colorectal carcinogenesis. Data were collected on the amounts of CD4+, CD8+ and CD56+ as well as on ThPOK+ cells infiltrated in normal colorectal mucosa (NM), in dysplastic aberrant crypt foci (microadenomas, MA, the earliest detectable lesions in colorectal carcinogenesis) and in colorectal carcinomas (CRC); moreover, the colocalization of ThPOK with the above-mentioned markers of immune cells was evaluated using confocal microscopy. Interestingly, ThPOK showed a prominent increase since MA. A strong colocalization of ThPOK with CD4 both in NM and in MA was observed, weaker in carcinomas. Surprisingly, there was a peak in the colocalization levels of ThPOK with CD8 in MA, which was evident, although to a lesser extent, also in carcinomas. In conclusion, according to the data of the present study, ThPOK may be considered a central regulator of the earliest events in the immune system during colorectal cancer development. The novelty of the present study is the proposed role of ThPOK in influencing the immune response against cancer cells.

References

- [1] Dunn et al. (2004) The immunobiology of cancer immunosurveillance and immunoediting. *Immunity* 21: 137-148.
- [2] Knaapen et al. (2006) Neutrophils and respiratory tract DNA damage and mutagenesis: a review. *Mutagenesis* 21: 225-236.
- [3] Coussens and Werb (2002) Inflammation and cancer. *Nature* 420: 860-867.
- [4] Watt and House (1978) Colonic carcinoma: a quantitative assessment of lymphocyte infiltration at the periphery of colonic tumors related to prognosis. *Cancer* 41: 279-282.
- [5] Galon et al. (2006) Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science* 313: 1960-1964.
- [6] Pagès et al. (2009) In situ cytotoxic and memory T cells predict outcome in patients with early-stage colorectal cancer. *J Clin Oncol* 27: 5944-5951.
- [7] Mlecnik et al. (2010) Biomolecular network reconstruction identifies T-cell homing factors associated with survival in colorectal. *Gastroenterology* 138: 1429-1434.

Keywords: Human colorectal cancer, immune response, ThPOK.