

This is the peer reviewed version of the following article:

Coffee and caffeine consumption in women affected by hypertension / Mattioli, Anna Vittoria; Ballerini Puviani, Matteo; Farinetti, Alberto. - In: JOURNAL OF HUMAN HYPERTENSION. - ISSN 0950-9240. - 32:6(2018), pp. 463-464. [10.1038/s41371-018-0048-z]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

01/05/2026 06:36

(Article begins on next page)



Coffee and caffeine consumption in women affected by hypertension

Anna Vittoria Mattioli¹ · Matteo Ballerini Puviani² · Alberto Farinetti¹

Received: 6 January 2018 / Revised: 30 January 2018 / Accepted: 7 February 2018
© Macmillan Publishers Limited, part of Springer Nature 2018

Dear Editor,

We have read with great interest the paper “Coffee consumption and risk of hypertension: a systematic review and dose–response meta-analysis of cohort studies” by Xie et al. [1] and we found their conclusion of importance with a view to clinical prevention. Authors concluded that their meta-analysis provides quantitative evidence that consumption of coffee was inversely associated with the risk of hypertension in a dose–response manner.

With reference to the findings reported in the paper, we would like to make the following contribution to the discussion. Recently, we evaluated 650 healthy women at premenopausal age, within this group we selected 237 women treated for hypertension (age range 45–54 years).

Women were categorized according to adherence to Mediterranean Diet, and group of food and beverages (fruit, vegetables, cereals, wine, and coffee) were singularly analysed. Caffeine intake was evaluated: One cup of espresso coffee included 90 mg of caffeine, one cappuccino = 110 mg of caffeine, one cup of American coffee = 160 mg, one can of cola soda = 42 mg, one can of energy drink = 80–120 mg, and one chocolate snack = 6 mg [2]. All women underwent evaluation of ABI index, a biomarker of pre-clinical atherosclerosis [3]. Local Ethics Institutional Board approved the protocol. Patients signed an informed consent. We found that coffee intake was higher in women in the high quartile of adherence to Mediterranean Diet. Hypertensive women that had high coffee consumption (>4 cups/day) had less frequently pre-clinical atherosclerosis

compared to women that consumed 1 cup/day or less [4]. We also found that the intake of chocolate snacks and bars increased caffeine intake from 150 ± 30 mg/day from coffee to 200 ± 50 mg/day from coffee plus chocolate.

Chocolate is a source of antioxidants leading to a beneficial effect. It is well known that the Mediterranean Diet is characterized by high intake of antioxidants, and, in our population, coffee and chocolate represents a source of antioxidants with an impact on total antioxidant intake [2, 3, 5, 7]. Coffee included several compounds and the bioavailability and the distribution of each compound and its metabolites contribute to coffee mechanisms of action. The mechanism of cardiovascular prevention is probably not related to caffeine but to one or more of the other compounds of coffee, i.e., chlorogenic acid, cafestol. Similarly, chocolate exerts beneficial effects on vessels [7]. A previous paper confirms this data in women, Rhee and coworkers found that caffeinated coffee, decaffeinated coffee, and caffeine are not risk factors for hypertension in postmenopausal women [8]. This could help to explain [9] the results from Xie et al. [1]. We can conclude that in our population of women living in the Mediterranean Area, the intake of coffee and caffeine do not increase significantly blood pressure. However what we found is merely an association and there are many confounders. The main confounder is the presence of the healthy “Mediterranean” diet and we cannot attribute the findings of our study to the caffeine intake alone.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest. Informed consent was obtained from all individual 66 participants included in the study.

References

1. Xie C, Cui L, Zhu J, Wang K, Sun N, Sun C. Coffee consumption and risk of hypertension: a systematic review and dose–response

✉ Anna Vittoria Mattioli
annavittoria.mattioli@unimore.it

¹ Department of Surgical, Medical and Dental Department of Morphological Sciences related to Transplant, Oncology and Regenerative Medicine, University of Modena and Reggio Emilia, Modena, Italy

² Istituto Nazionale per le Ricerche Cardiovascolari U.O. di Modena, Modena, Italy

- 71 meta-analysis of cohort studies. *J Hum Hypertens*. 2018. <https://doi.org/10.1038/s41371-017-0007-0>
- 72
- 73 2. Mattioli AV. Effects of caffeine and coffee consumption on cardiovascular disease and risk factors. *Future Cardiol*. 2007;3:203–12.
- 74
- 75
- 76 3. Mattioli AV, Coppi F, Migaldi M, Scicchitano P, Ciccone MM, Farinetti A. Relationship between Mediterranean diet and asymptomatic peripheral arterial disease in a population of premenopausal women. *Nutr Metab Cardiovasc Dis*. 2017;27:985–90.
- 77
- 78
- 79
- 80 4. Mattioli AV, Migaldi M, Farinetti A. Coffee in hypertensive women with asymptomatic peripheral arterial disease: a potential nutraceutical effect. *J Cardiovasc Med*. 2018. in press
- 81
- 82
- 83
- 84 5. Mattioli AV, Farinetti A, Miloro C, Pedrazzi P, Mattioli G. Influence of coffee and caffeine consumption on atrial fibrillation in hypertensive patients. *Nutr Metab Cardiovasc Dis*. 2010;21:412–7.
- 85
- 86 6. D’Elia L, La Fata E, Galletti F, Scalfi L, Strazzullo P. Coffee consumption and risk of hypertension: a dose–response meta-analysis of prospective studies. *Eur J Nutr*. 2017. <https://doi.org/10.1007/s00394-017-1591-z>
- 87
- 88
- 89
- 90 7. Mattioli AV, Miloro C, Pennella S, Pedrazzi P, Farinetti A. Adherence to Mediterranean diet and intake of antioxidants influence spontaneous conversion of atrial fibrillation. *Nutr Metab Cardiovasc Dis*. 2013;23:115–121.
- 91
- 92
- 93
- 94 8. Noordzij M, Uiterwaal CS, Arends LR, Kok FJ, Grobbee DE, Geleijnse JM. Blood pressure response to chronic intake of coffee and caffeine: a meta-analysis of randomized controlled trials. *J Hypertens*. 2005;23:921–8.
- 95
- 96
- 97
- 98 9. Rhee JJ, Qin F, Hedlin HK, Chang TI, Bird CE, Zaslavsky O, et al. Coffee and caffeine consumption and the risk of hypertension in postmenopausal women. *Am J Clin Nutr*. 2016;103:210–07.
- 99
- 100

UNCORRECTED PROOF

Journal : 41371

Article : 48

SPRINGER NATURE

Author Query Form

Please ensure you fill out your response to the queries raised below and return this form along with your corrections

Dear Author

During the process of typesetting your article, the following queries have arisen. Please check your typeset proof carefully against the queries listed below and mark the necessary changes either directly on the proof/online grid or in the 'Author's response' area provided below

Queries	Details Required	Author's Response
AQ1	Author surnames have been highlighted - please check these carefully and indicate if the first name or surname have been marked up incorrectly. Please note that this will affect indexing of your article, such as in PubMed.	
AQ2	Should you wish to order offprints, please click on www.nature.com/aj/forms/jhh_offprint_2017.pdf to download and complete the offprint form and upload the completed form along with the article.	
AQ3	Please clarify whether the splitting of the original affiliation 1,2 as done is correct. If No, please show how they should be sorted out.	
AQ4	Please note that reference [6] included in the reference list, but has not cited in text part. Please supply details for the missing reference.	
AQ5	Please update ref. 4.	
AQ6	Please confirm edits made in conflict of interest is ok.	