This is the peer reviewd version of the followng article:

O-312 Occupational exposure levels of solar ultraviolet radiation: an EPHOR review / Würtz, E.T., Cherrie, M., Ge, C., Grandahl, K., Hansen, J., Dahlman-Höglund, A., Mehlum, I.S., Modenese, A., Notö, H., Peters, C., Selander, J., Solovieva, S., Schlünssen, V., Wittlich, M., Kolstad, H.A.. - In: OCCUPATIONAL AND ENVIRONMENTAL MEDICINE. - ISSN 1470-7926. - 78:Suppl. 1(2021), pp. A46.2-A46. (28th International Symposium on Epidemiology in Occupational Health "From the Workplace to the Population: Exposure and Prevention" - EPICOH 2021 Online 25-28 October 2021) [10.1136/OEM-2021-EPI.123].

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

13/11/2026 00:11

(Article begins on next page)

Occupational exposure levels of solar ultra violet radiation: an EPHOR review

*Else Toft Würtz[1], Mark Cherrie[2], Calvin Ge[3], Kasper Grandahl[4], Johnni Hansen[5], Anna Dahlman-Hoglund[6], Ingrid Sivesind Mehlum[7], Alberto Modenese[8], Hilde Notö[7], Cheryl Peters[9], Jenny Selander[10], Svetlana Solvieva[11], Vivi Schlünssen[12], Marc Wittlich[13], Henrik Albert Kolstad[1]

[1]Department of Occupational Medicine, Aarhus University Hospital, Denmark
[2]Institute of Occupational Medicine, Edinburgh University, Edinburgh, UK
[3]Institute for Risk Assessment Sciences (IRAS), Utrecht University, Utrecht, Netherlands
[4]Department of Occupational Medicine, Copenhagen University Holbaek, Holbaek, Denmark
[5]The Danish Cancer Society Research Center, Copenhagen, Denmark
[6]Department of Occupational and Environmental Medicine, Sahlgrenska University Hospital, University of Göteborg, Göteborg, Sweden
[7]National Institute of Occupational Health (STAMI), Oslo, Norway
[8]Occupational Medicine, Department of Biomedical, Metabolic and Neural Sciences, University of Modena & Reggio Emilia, Modena, Italy
[9]Cancer Epidemiology and Prevention Research, Alberta Health Services, Calgary, AB, Canada
[10]Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden
[11]Finnish Institute of Occupational Health, Helsinki, Finland
[12]Department of Public Health, Aarhus University, Aarhus, Denmark
[13]Institute for Occupational Safety and Health of the German Social Accident Insurance, Sankt Augustin, Germany

Introduction: Solar ultra violet radiation (UVR) causes malignant melanoma, squamous cell carcinoma, and basal cell carcinoma of the skin, is a risk factor of cataract, and the primary predictor of vitamin-D levels. The UVR exposure level depends on several factors such as time of the day, season, activity level, latitude and altitude, and vary across occupations, with high levels reported in different outdoor jobs. Personal measurements are typically reported as full shift cumulative Standard Erythema Dose (SED).

Objective: To conduct a systematic review of occupational solar UVR exposure levels.

Methods: We performed a PubMed search for literature presenting personal solar UVR levels measured by wearable UVR sensors during work and supplemented with papers identified from reference lists. We screened title and abstract of 1005 papers, full text reviewed 113 papers, and extracted data from 42 included papers. Job descriptions were coded according to the International Standard Classification of Occupations (ISCO-88) on the 4-digit level.

Results: In total, UV exposure levels of 35345 personal workday measurements of mainly construction workers, farmers, seamen, teachers, expeditioners, and athletes were included. They covered all year, 6 am - 6 pm work hours, latitudes from 20-75° north and 10-68° south, and altitudes in the 0-8000 m range. 76.2% of the studies included repeated measurements. Eleven different polysulphone film, UV-sensitive spore, or electronic irradiance sensors placed on head, neck, shoulder, chest, back, arm, or wrist were used. Exposure levels varied between 0.4 SED for laboratory workers and 28.6 SED for construction workers.

Conclusion: The established measurement database will be used for the establishment of a general population quantitative job exposure matrix (JEM) as part of the European Exposome Project for Health and Occupational Research (EPHOR). The UV-JEM will be applied in multi-centre European occupational cohort studies of adverse and gainful effects of exposure to UVR.