

This is the peer reviewed version of the following article:

Riluzole and other prognostic factors in ALS: a population-based registry study in Italy / Mandrioli, Jessica; Malerba, Sara Angela; Beghi, Ettore; Fini, Nicola; Fasano, Antonio; Zucchi, Elisabetta; De Pasqua, Silvia; Guidi, Carlo; Terlizzi, Emilio; Sette, Elisabetta; Ravasio, Alessandro; Casmiro, Mario; Salvi, Fabrizio; Liguori, Rocco; Zinno, Lucia; Handouk, Yasmin; Rizzi, Romana; Borghi, Annamaria; Rinaldi, Rita; Medici, Dorian; Santangelo, Mario; Granieri, Enrico; Mussuto, Vittoria; Aiello, Marina; Ferro, Salvatore; Vinceti, Marco. - In: JOURNAL OF NEUROLOGY. - ISSN 0340-5354. - 265:4(2018), pp. 817-827. [10.1007/s00415-018-8778-y]

Terms of use:

The terms and conditions for the reuse of this version of the manuscript are specified in the publishing policy. For all terms of use and more information see the publisher's website.

27/04/2026 01:08

(Article begins on next page)

27/04/2026 01:08

Journal of Neurology

Riluzole and other prognostic factors in ALS: a population-based registry study in Italy

--Manuscript Draft--

Manuscript Number:	JOON-D-17-01262R3
Full Title:	Riluzole and other prognostic factors in ALS: a population-based registry study in Italy
Article Type:	Original Communication
Corresponding Author:	JESSICA MANDRIOLI, MD St. Agostino Estense Hospital Modena, Modena ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	St. Agostino Estense Hospital
Corresponding Author's Secondary Institution:	
Corresponding Author E-Mail:	j.mandrioli@ausl.mo.it
First Author:	JESSICA MANDRIOLI, MD
First Author Secondary Information:	
Order of Authors:	JESSICA MANDRIOLI, MD
	Sara Angela Malerba
	Ettore Beghi
	Nicola Fini
	Antonio Fasano
	Elisabetta Zucchi
	Silvia De Pasqua
	Carlo Guidi
	Emilio Terlizzi
	Elisabetta Sette
	Alessandro Ravasio
	Mario Casmiro
	Fabrizio Salvi
	Rocco Liguori
	Lucia Zinno
	Yasmin Handouk
	Romana Rizzi
	Annamaria Borghi
	Rita Rinaldi
	Doriana Medici
	Mario Santangelo
	Enrico Granieri
	Vittoria Mussuto
	Marina Aiello

	Salvatore Ferro	
	Marco Vinceti	
Order of Authors Secondary Information:		
Funding Information:	Emilia Romagna Registry for ALS is supported by a grant from the Emilia Romagna Regional Health Authority.	Dr. JESSICA MANDRIOLI
Abstract:	<p>Objective: In this prospective population-based registry study on ALS survival, we investigated the role of riluzole treatment, together with other clinical factors, on the prognosis in incident ALS cases in Emilia Romagna Region (ERR), Italy.</p> <p>Methods: A registry for ALS has been collecting all incident cases in ERR since 2009. Detailed clinical data from all patients diagnosed with ALS between 1.1.2009 and 31.12.2014 have been analyzed for this study, with last follow up date set at 31.12.2015.</p> <p>Results: During the 6 years of the study, there were 681 incident cases with a median tracheostomy-free survival of 40 months (95%C.I. 36-44) from onset and of 26 months (95%C.I. 24-30) from diagnosis; 573 patients (84.14%) were treated with riluzole, 207 (30.39%) patients underwent gastrostomy, 246 (36.12%) non invasive ventilation, and 103 (15.15%) invasive ventilation. Patients who took treatment for $\geq 75\%$ of disease duration from diagnosis had a median survival of 29 months compared to 18 months in patients with $< 75\%$ treatment duration. In multivariable analysis, factors independently influencing survival were age at onset (HR 1.04, 95%CI 1.02-1.05, $p < 0.001$), dementia (HR 1.56, 95%CI 1.05-2.32, $p = 0.027$), degree of diagnostic certainty (HR 0.88, 95%CI 0.78-0.98, $p = 0.021$), gastrostomy (HR 1.46, 95%CI 1.14-1.88, $p = 0.003$), NIV (HR 1.43, 95%CI 1.12-1.82, $p = 0.004$), and weight loss at diagnosis (HR 1.05, 95%CI 1.03-1.07, $p < 0.001$), diagnostic delay (HR 0.98, 95%CI 0.97-0.99, $p = 0.004$), and % treatment duration (HR 0.98, 95%CI 0.98-0.99, $p < 0.001$).</p> <p>Conclusions: Independently from other prognostic factors, patients who received riluzole for a longer period of time survived longer, but further population based studies are needed to verify if long-term use of riluzole prolongs survival.</p>	
Response to Reviewers:	<p>Response to reviewer comments on "Riluzole and other prognostic factors in ALS: a population-based registry study in Italy"</p> <p>We wish to thank the reviewers and editor for their comprehensive review of our manuscript. We have done our best to address each concern. We are grateful to Journal of Neurology for the opportunity to submit the revised manuscript.</p> <p>Best regards, Jessica Mandrioli (on behalf of all coauthors)</p> <p>Reviewer: 1</p> <p>Comments for the Author:</p> <p>Reviewer #1: COMMENTS FOR AUTHOR:</p> <p>Reviewer #1:</p> <p>1) Survival is defined as "diagnosis to death" (page 6, line 40) but time-varying covariate is used for "time from onset to riluzole exposure" (page 7, line 10-14 and page 8, line 34-36). The time-varying covariate should be in accordance with the survival period definition. Thus, if survival is used "from diagnosis to death" then the time-varying covariate should also be used "from diagnosis" to riluzole exposure (and not from onset).</p> <p>We agree and we made the suggested changes (page 7, lines 10-12 and page 8, lines 34-36) Results were corrected accordingly (page 8 lines 38-45, 54-61; page 9 lines 4-6, 16-26, table 4)</p> <p>2) Figure 1a: Crossing of curves at about 24 months with the riluzole group associated with a rather detrimental effect after 84 months. This is an interesting finding also described in other studies. The authors may wish to discuss the discrepancy between</p>	

	<p>the lack of an overall riluzole effect (figure 1a) and the remarkably beneficial effect in the other analysis, i.e. % treatment duration with HR = 0.18.</p> <p>We discuss this finding as suggested (discussion page 10 lines 12-24)</p> <p>3) Figure 1c: Should be PDC >90</p> <p>We changed it accordingly.</p> <p>4) "% treatment duration (HR 0.18...)" (on page 4, line 32): This seems to be the main result but is not mentioned in the result section (it is only listed in table 4 on the last page). Moreover, such a HR is extremely low and would mean that each % increase in treatment duration would reduce ALS mortality to only 18%. The authors may wish to double-check their calculations and analysis.</p> <p>We apologize for the mistake, due to the fact that we calculated PDC as the ratio between Riluzole treatment duration/disease duration and not as % (as a consequence the HR of 0.18 refers to an increase of PDC of 1 unit, that is not possible). Considering PDC as %, after correcting for immortal time bias, we obtained a HR of 0.98, i.e. a decrease of the risk of death/tracheostomy of 2% for each % increase in treatment duration.</p> <p>We added the main result in the results section (page 8, lines 38-45). We corrected the mistake also on page 8 lines 58-60, on page 9 lines 4-6 and 16-19 and in table 2 and table 4.</p> <p>—</p>
<p>Author Comments:</p>	<p>Modena, 23rd January 2018</p> <p>Dear Sir,</p> <p>we would like to submit our revised manuscript entitled "Riluzole and other prognostic factors in ALS: a population-based registry study in Emilia Romagna, Italy" for consideration by Journal of Neurology, as an original communication.</p> <p>We performed a prospective population-based registry study on ALS survival, aiming at studying the role of Riluzole treatment, together with other clinical factors, on the prognosis in incident ALS cases in Emilia Romagna Region (ERR), Italy. A registry for ALS has been collecting all incident cases in ERR since 2009. Detailed clinical data from all patients diagnosed with ALS between 1.1.2009 and 31.12.2014 have been analyzed for this study, with last follow up date set at 31.12.2015. During the 6 years of the study, there were 681 incident cases with a median tracheostomy-free survival of 40 months (95%C.I. 36-44) from onset and of 26 months (95%C.I. 24-30) from diagnosis; 573 patients (84.14%) were treated with riluzole, 207 (30.39%) patients underwent gastrostomy, 246 (36.12%) non invasive ventilation, and 103 (15.15%) invasive ventilation. Patients who took treatment for ≥75% of disease duration had a median survival of 29 months compared to 18 months in patients with <75% treatment duration. In multivariable analysis, factors independently influencing survival were age at onset (HR 1.04,95%CI 1.02-1.05,p<0.001), dementia (HR 1.56,95%CI 1.05-2.32, p=0.027), degree of diagnostic certainty (HR 0.88,95%CI 0.78-0.98,p=0.021), gastrostomy (HR1.46, 95%CI 1.14-1.88,p=0.003), NIV (HR1.43,95%CI 1.12-1.82,p=0.004), and weight loss at diagnosis (HR1.05,95%CI1.03-1.07,p<0.001), diagnostic delay (HR 0.98, 95%CI 0.97-0.99,p=0.004), and % treatment duration (HR 0.98, 95%CI 0.98-0.99,p<0.001).</p> <p>In summary, in our study, independently from other prognostic factors, patients who received riluzole for a longer period of time survived longer, but further population based studies are needed to verify if long-tem use of riluzole prolongs survival.</p> <p>We hope to have accomplished with all the reviewers requests. We hope that you will find our work to be of interest to Journal of Neurology readership and to the scientific community. The authors declare that they do not have any conflict of interest. All the authors have read the final draft of the manuscript and declare that there isn't any other author not included in the list. The manuscript has not been previously published and it is not under simultaneous consideration by other journals.</p>

Yours sincerely,

Dr. Jessica Mandrioli
Department of Neurosciences, Nuovo Ospedale Civile S. Agostino Estense
Via P.Giardini 1355, 41126 Modena,Italy
Tel. +390593961640, Fax +390593963775
Cell +393392633246
E-Mail j.mandrioli@ausl.mo.it

TITLE PAGE**Authors:**

Jessica Mandrioli¹, MD, Sara Angela Malerba¹, MD, Ettore Beghi², MD, Nicola Fini¹, MD, Antonio Fasano¹, MD, Elisabetta Zucchi¹, MD, Silvia De Pasqua³, MD, Carlo Guidi⁴, MD, Emilio Terlizzi⁵, MD, Elisabetta Sette⁶, MD, Alessandro Ravasio⁷, MD, Mario Casmiro⁸, MD, Fabrizio Salvi⁹, MD, Rocco Liguori³, MD, Lucia Zinno¹⁰, MD, Yasmin Handouk¹¹, Romana Rizzi¹², MD, Annamaria Borghi¹³, MD, PhD, Rita Rinaldi¹⁴, MD, PhD, Doriana Medici¹⁵, MD, Mario Santangelo¹⁶, MD, Enrico Granieri¹⁷, MD, Vittoria Mussuto¹⁸, MD, Marina Aiello¹⁹, MD, Salvatore Ferro²⁰, ERRALS Group, and Marco Vinceti²¹, MD, PhD.

Title:**Riluzole and other prognostic factors in ALS: a population-based registry study in Italy**

- (1) Department of Neuroscience, St. Agostino-Estense Hospital, Azienda Ospedaliero Universitaria di Modena, University of Modena and Reggio Emilia, Modena, Italy
- (2) Department of Neuroscience, IRCCS-Istituto di Ricerche Farmacologiche Mario Negri, Milano, Italy
- (3) Department of Biomedical and Neuromotor Sciences, University of Bologna and IRCCS Istituto delle Scienze Neurologiche di Bologna, Bologna, Italy
- (4) Department of Neurology, Forlì Hospital, Forlì, Italy
- (5) Department of Neurology, G. Da Saliceto Hospital, Piacenza, Italy
- (6) Department of Neurology, St. Anna Hospital, Ferrara, Italy
- (7) Department of Neurology, Infermi Hospital, Rimini, Italy
- (8) Department of Neurology, Faenza and Ravenna Hospital, Ravenna, Italy
- (9) Neurology Unit, Bellaria Hospital, IRCCS Istituto delle Scienze Neurologiche di Bologna, Bologna, Italy
- (10) Department of Neuroscience, Neurology Unit, University of Parma, Parma, Italy
- (11) Department of Neurology, Bufalini Hospital, Cesena, Italy
- (12) Department of Neurology, IRCCS Arcispedale Santa Maria Nuova, Reggio Emilia, Italy
- (13) Neurology Unit, Maggiore Hospital, IRCCS Istituto delle Scienze Neurologiche di Bologna, Bologna, Italy
- (14) Department of Neuroscience, St. Orsola-Malpighi University Hospital, Bologna, Italy
- (15) Department of Neurology, Fidenza Hospital, Parma, Italy
- (16) Department of Neurology, Carpi Hospital, Modena, Italy
- (17) Department of Neuroscience, University of Ferrara, Ferrara, Italy

(18) Department of Neurology, Imola Hospital, Bologna, Italy

(19) Department of Clinical and Experimental Medicine, Respiratory Disease and Lung Function Unit, University of Parma, Parma, Italy.

(20) Department of Hospital Services, Emilia Romagna Regional Health Authority, Bologna, Italy

(21) Environmental, Genetic and Nutritional Epidemiology Research Center (CREAGEN), University of Modena and Reggio Emilia Medical School, Modena, Italy;

Corresponding author:

Jessica Mandrioli,MD

Department of Neuroscience, St. Agostino-Estense Hospital,Azienda Ospedaliero Universitaria di Modena

Via Pietro Giardini n. 1355

41100 Modena, Italy

Tel. 00390593961640 Fax 00390593963775

Cell +303392633246

E-Mail j.mandrioli@ausl.mo.it

DISCLOSURE OF CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest

ACKNOWLEDGEMENTS

The authors acknowledge ERRALS group:

Project coordinator: J. Mandrioli

Collaborating centers:

Department of Neuroscience, St. Agostino Estense Hospital, Azienda Ospedaliero Universitaria di Modena, Italy (J.

Mandrioli, N. Fini, A. Fasano, E. Zucchi, A. Gessani, L. Ferri, S. Malerba, P. Nichelli)

Department of Neurology, Bufalini Hospital, Cesena, Italy (Y. Handouk, S. Biguzzi, W. Neri)

Department of Neurology, Forlì Hospital, Forlì, Italy (C. Guidi, W. Neri)

Department of Neurosciences and Rehabilitation, St Anna Hospital, Ferrara (E. Sette, V. Tugnoli and M.R. Tola)

Department of Neurology, G. Da Saliceto Hospital, Piacenza, Italy (E. Terlizzi, D. Guidetti)

Department of Neurology, Infermi Hospital, Rimini, Italy (M. Currò Dossi, M. Pasquinelli, J. Andruccioli, A. Ravasio)

Department of Neurology, Faenza and Ravenna Hospital, Ravenna, Italy (M. Casmiro,F. Rasi)

Department of Neurology and IRCCS Istituto delle Scienze Neurologiche di Bologna , Bellaria Hospital, Bologna, Italy

(F. Salvi, I. Bartolomei, R. Michelucci)

1 Department of Biomedical and Neuromotor Science and IRCCS Istituto delle Scienze Neurologiche di Bologna,
2 University of Bologna, Bologna, Italy (P. Avoni, S. De Pasqua, R. Liguori)

3
4 Department of Neurology, IRCCS Arcispedale Santa Maria Nuova, Reggio Emilia, Italy (R. Rizzi, E. Canali, L.
5 Codeluppi, F. Valzania)

6
7 Department of Neuroscience, University of Parma, Parma, Italy (A. Taga, V. Pietrini, M.C. Trapasso, and L. Zinno)

8
9 Department of Clinical and Experimental Medicine, Respiratory Disease and Lung Function Unit, University of Parma,
10 Parma, Italy (M. Aiello)

11
12 Department of Neurology, Fidenza Hospital, Parma, Italy (D. Medici, E. Chierici, E. Montanari)

13
14 Department of Neurology, Carpi Hospital, Modena, Italy (M. Santangelo, S. Amidei)

15
16 Department of Biomedical and Specialty Surgical Sciences,, University of Ferrara, Ferrara, Italy (I. Casetta, M.
17 Pugliatti, E. Della Coletta, V. Govoni, E. Granieri)

18
19 Department of Neurology, Imola Hospital, Bologna, Italy (P. De Massis, V. Mussuto)

20
21 Department of Neurology and IRCCS Istituto delle Scienze Neurologiche di Bologna, Maggiore Hospital, Bologna,
22 Italy (A. Borghi, A. Gabellini, R. Michelucci)

23
24 Neurology Unit, St. Orsola-Malpighi University Hospital, AOU di Bologna, Italy (R. Rinaldi, F. Cirignotta)

25
26 Department of Hospital Services, Emilia Romagna Regional Health Authority, Bologna, Italy (S. Ferro)

27
28 Neuroepidemiology Unit, IRCCS Istituto delle Scienze Neurologiche di Bologna, Italy (R. D'Alessandro)

29
30 CREAGEN Research Center for Environmental, Genetic and Nutritional Epidemiology, University of Modena and
31 Reggio Emilia, Modena, Italy

32 33 34 35 36 37 38 39 40 **FUNDING**

41
42 Emilia Romagna Registry for ALS is supported by a grant from the Emilia Romagna Regional Health Authority.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

ABSTRACT

1
2
3
4 **Objective:** In this prospective population-based registry study on ALS survival, we investigated the role of riluzole
5 treatment, together with other clinical factors, on the prognosis in incident ALS cases in Emilia Romagna Region
6 (ERR), Italy.
7

8
9
10 **Methods:** A registry for ALS has been collecting all incident cases in ERR since 2009. Detailed clinical data from all
11 patients diagnosed with ALS between 1.1.2009 and 31.12.2014 have been analyzed for this study, with last follow up
12 date set at 31.12.2015.
13
14

15
16 **Results:** During the 6 years of the study, there were 681 incident cases with a median tracheostomy-free survival of 40
17 months (95% C.I. 36-44) from onset and of 26 months (95% C.I. 24-30) from diagnosis; 573 patients (84.14%) were
18 treated with riluzole, 207 (30.39%) patients underwent gastrostomy, 246 (36.12%) non invasive ventilation, and 103
19 (15.15%) invasive ventilation. Patients who took treatment for $\geq 75\%$ of disease duration from diagnosis had a median
20 survival of 29 months compared to 18 months in patients with $< 75\%$ treatment duration. **In multivariable analysis,**

21
22 **factors independently influencing survival were age at onset (HR 1.04, 95% CI 1.02-1.05, $p < 0.001$), dementia (HR**
23 **1.56, 95% CI 1.05-2.32, $p = 0.027$), degree of diagnostic certainty (HR 0.88, 95% CI 0.78-0.98, $p = 0.021$), gastrostomy**
24 **(HR 1.46, 95% CI 1.14-1.88, $p = 0.003$), NIV (HR 1.43, 95% CI 1.12-1.82, $p = 0.004$), and weight loss at diagnosis**
25 **(HR 1.05, 95% CI 1.03-1.07, $p < 0.001$), diagnostic delay (HR 0.98, 95% CI 0.97-0.99, $p = 0.004$), and % treatment duration**
26 **(HR 0.98, 95% CI 0.98-0.99, $p < 0.001$).**
27
28
29
30
31
32
33
34
35

36 **Conclusions:** Independently from other prognostic factors, patients who received riluzole for a longer period of time
37 survived longer, but further population based studies are needed to verify if long-term use of riluzole prolongs survival.
38
39
40

41
42 **Key words:** amyotrophic lateral sclerosis, survival, prognostic factors, therapeutic intervention, Riluzole.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

TEXT

INTRODUCTION

Amyotrophic lateral sclerosis (ALS) is a rare neurodegenerative disease characterized by progressive motor deficits over the course of weeks to months leading to severe disability and death. The disease is highly heterogeneous in clinical presentation, age at onset, involvement of other than motor systems (cognitive impairment in up to 50%, variable sensory, autonomic, extrapyramidal signs), family history and genetic background, disease course and survival [1, 2]. Although survival of ALS patients from symptom onset is reported to be 3-5 years, published studies report a wide range of outcomes influenced by age and site of onset, severity and rate of disease progression, diagnostic delay, cognitive impairment, nutritional and respiratory status, functional/disability scores, Revised El Escorial diagnostic Criteria (EEC-R), multidisciplinary approach, and therapeutic interventions [3–5].

Despite significant efforts and a large number of well-designed clinical trials, published studies on ALS treatments gave negative results, probably because of ALS phenotypic, genetic and pathophysiological heterogeneity [6]. So far, riluzole is the only drug that has been shown to increase survival in ALS patients [7], although with a poorly understood mechanism of action[8] and a way of drug delivery still investigational [9, 10].

Riluzole has been tested in four randomized clinical trials (RCTs) involving 1477 patients, showing a significant effect on survival which was quantified as a gain of 3 months in 50% of patients or a 9% gain in the probability of surviving one year [11].

Conversely, subsequent cohort studies have suggested that treatment with riluzole may be associated with a median survival increase of 6 months to as many as 12 months, with diverse impact on different subpopulations of ALS patients [5, 12–15]. Two recent studies (one retrospective [16] and one prospective [17]) considered the cumulative defined daily dose to measure treatment duration, and obtained contrasting results.

In this prospective population-based registry study on ALS survival, our aim was to study the role of riluzole treatment and its duration on prognosis in incident ALS cases in Emilia Romagna Region (ERR), Italy, between 2009 and 2014.

METHODS

Patient data collection

The study was performed in Emilia Romagna Region (ERR), northern Italy, considering patients diagnosed with ALS from 1/1/2009 to 31/12/2014. The region of Emilia-Romagna covers an area of 22,453 km², ranking 6th in Italy, and during the evaluated time period, the local population increased from 4,395,569 (2009) to 4,450,508 (2014).

A prospective registry (Emilia Romagna Registry for ALS–ERRALS) has been active in our region since 2009 [18], collecting all incident ALS cases among residents of ERR, diagnosed according to EEC-R [19]. Caring physicians

collected a detailed phenotypic profile of each ALS patient, including the following parameters: age at onset and diagnosis, gender, type of onset, site and time of onset, affected regions, upper and lower motor neuron signs, EEC-R classification, clinical phenotype [1], presence of dementia (diagnosed by neuropsychological testing) and/or extrapyramidal signs, family history, and drugs (including riluzole).

These data have been included at diagnosis into an electronic database available through a dedicated web-site, accessible only to the investigators of ERR Neurological Departments. In each department, one or more investigators were identified as study referents, and had to upload data on new ALS cases as soon as possible after diagnosis. Clinical follow up has been performed in the 17 neurological departments of ERR until death, collecting and uploading information on ALS course, including forced vital capacity (FVC), ALS Functional Rating Scale – Revised (ALSFRS-R) [20, 21], riluzole intake and discontinuation, gastrostomy [22], non-invasive (NIV) or invasive (IV) ventilation support, and cause, place and time of death. A regular supervision on the data has been performed by the coordinating center, checking for accuracy and completeness.

Conforming to the EFNS Guidelines [23], patients underwent a regular multidisciplinary follow-up at least every 3-4 months. When patients were no longer able to reach the centers, they were monitored at their home or in a nursing home by the Integrated Home Care of the Italian National Health System (INHS)[24].

For this study we focused on clinical variables and riluzole intake, considering date of riluzole administration and discontinuation (if applicable) with respect to disease duration.

The study was approved by the ethical committees of the coordinating center and of ERR provinces.

Statistical Methods

Continuous variables are reported as mean and standard deviation (SD) from the mean. Discrete variables are reported as number and proportion of subjects with the characteristic of interest. Descriptive statistics were performed using Student's t-test and chi-square test where appropriate. Survival was calculated from onset and from diagnosis to death/tracheostomy or the censoring date (last day of follow-up, 31/12/2015) using the Kaplan-Meier method. The curves were compared with the Log-Rank test. Multivariable analysis was performed with Cox's proportional hazards model, to evaluate the association between demographic and clinical factors, riluzole use/treatment duration and survival.

Riluzole use was defined as a categorical variable (0='no drug', 1='drug user' if treatment was taken at a dosage of 100 mg for at least 1 month), but we also classified individuals in different groups based on the duration of riluzole treatment (in days from treatment initiation to treatment end or death/tracheostomy) with respect to ALS duration (in days from diagnosis to death/tracheostomy) (proportion of days covered -PDC)[16]. PDC definition was adopted because an evaluation of survival times exclusively in dependence of the days of riluzole treatment would have biased

the results since patients with longer therapy durations will have lived necessarily longer [16]. Mean PDC was 75%, median PDC was 90%; we considered these values as cut off values for the individuation of three treatment groups. We also calculated the delay from onset of symptoms to start of riluzole (months). Regarding the analysis of riluzole's effect on survival we accounted for a possible bias due to a delay in the start of riluzole treatment after diagnosis by adjusting for the immortal time bias, which refers, in observational studies, to the time during which the outcome could not have occurred [25]. To correct for this bias we used a time-varying covariate **for time from diagnosis to Riluzole exposure** in order to avoid misclassification of exposed patients' survival time before the first prescription as the exposed follow-up time [25].

We measured ALSFRS-R at diagnosis and ALSFRS-R monthly decline, subtracting ALSFRS-R total score at diagnosis from the score of 48 presumed at onset and dividing the results for the number of months of diagnostic delay [26].

For all Cox proportional hazard regression analyses, we considered age at onset, phenotypes, type of onset, EEC-R classification, presence of dementia and/or extrapyramidal signs, comorbidities using the Charlson Comorbidity Index (CCI) [27], family history of ALS, monthly ALSFRS-R decline at diagnosis and riluzole use, to assess the independent contribution of the variables of interest to death or tracheostomy.

A p value <0.05 was considered significant. All calculations were performed with STATA statistical package, V.12 (2013).

RESULTS

Clinical characteristics of ALS patients

During the 6 years of study, there were 681, 371 men and 310 women (men:women =1.20), incident cases in ERR.

Table 1 shows the clinical features of ALS patients. Median survival time was 40 months (95% C.I. 36-44) from onset and 26 (95% C.I. 24-30) from diagnosis. The overall 1-year, 2-year, 3-year and 5-year survival rates from onset were 89.6%, 68.8%, 53.8% and 36.6% respectively. The overall 1-year, 2-year, 3-year and 5-year survival rates from diagnosis were 73.3%, 52.8%, 39.9% and 32.7% respectively. Five hundred and seventy-three patients (84.14%) were treated with riluzole, 207 (30.39%) underwent gastrostomy, 246 (36.12%) NIV, and 103 (15.15%) IV.

Univariate analysis of factors associated with ALS survival

According to univariate analysis, factors related to tracheostomy-free survival from diagnosis were (table 2): age at onset, diagnostic delay, site of onset, phenotype, presence of dementia, degree of diagnostic certainty according to EEC-R, weight change between healthy status and diagnosis, ALSFRS-R score at diagnosis, and ALSFRS-R monthly decline, NIV and gastrostomy, comorbidities (CCI). The same factors influenced survival from disease onset (table 2). Overall riluzole treatment did not influence ALS survival (figure 1A). In contrast, the duration of treatment in relation to disease duration, expressed as percent of days of riluzole use (until discontinuation, tracheostomy or death)(PDC),

influenced survival (Table 2). Standard daily dose of riluzole was 50 mg twice a day. Median PDC was 90%. Mean PDC was 75%. Patients that took riluzole for a number of days corresponding to at least 75% of disease duration from diagnosis ($\geq 75\%$ PDC) had a median survival of 29 months as compared to 18 months in patients with a shorter percentage of riluzole treatment (figure 1B). Patients that took riluzole for a number of days corresponding to at least 90% of disease duration from diagnosis ($\geq 90\%$ PDC) had a median survival of 46 months as compared to 15 months in patients with a shorter percentage of riluzole treatment (figure 1C). The delay from disease onset to start of treatment significantly influenced survival (table 2) as the earlier the patients took riluzole, the worse the prognosis. Half of treated patients took riluzole treatment after 317 days (~10 months) from onset.

Patients who took riluzole for $\geq 90\%$ PDC were younger, had a shorter diagnostic delay, a lower CCI and ALSFRS-R decline, had more frequently a spinal onset, were less frequently affected by dementia and underwent more often NIV and IV (table 3). There was a mean difference of 2 months in the delay from diagnosis to treatment initiation between the two groups.

Multivariable analysis of factors associated with ALS survival

We then did a multivariable analysis including PDC, the delay from onset to start of treatment, and all variables possibly influencing survival available at time of diagnosis. In the initial Cox multivariable model, we included the following variables (retention criterion of <0.10): diagnostic delay, age at onset, site of onset, phenotypes, presence of dementia, EEC-R classification, PDC, delay from onset to start of treatment, ALSFRS-R score at diagnosis, ALSFRS-R monthly decline at diagnosis, weight loss at diagnosis, comorbidities (CCI), gastrostomy and NIV. We also used a time-varying covariate for time from diagnosis to riluzole exposure.

After dropping non-significant terms (stepwise backward), the final model included age at onset, diagnostic delay, dementia, EEC-R classification, gastrostomy, NIV, weight loss at diagnosis, PDC (table 4). As for PDC (%), after correcting for immortal time bias, we obtained a HR of 0.98, i.e. a decrease of the risk of death/tracheostomy of 2% for each % increase in treatment duration.

Subgroups analysis of survival

Hypothesizing that early treatment discontinuation was most frequent in more severely affected patients, we excluded from our population those patients who died before 6 months from diagnosis (79 cases). In the remaining 602 patients the results were unchanged and factors independently influencing survival (multivariable analysis) were age at onset (HR 1.03, 95% C.I. 1.02-1.04, $p<0.001$), diagnostic delay (HR 0.97, 95% C.I. 0.96-0.99, $p=0.001$), EEC-R classification (HR 0.86, 95% C.I. 0.76-0.98, $p=0.020$), NIV (HR 1.71, 95% C.I. 1.31-2.24, $p<0.001$), gastrostomy (HR 1.61, 95% C.I. 1.23-2.11, $p=0.001$), weight loss at diagnosis (HR 1.04, 95% C.I. 1.01-1.06, $p=0.002$), and PDC (HR 0.98, 95% C.I. 0.98-0.99, $p=0.001$). As compared to the remaining study population, the 79 patients dying early were older (mean age

at onset 73.78 ± 7.89 years vs 65.53 ± 11.10 years), with bulbar onset (55.70% vs. 44.30%), more frequently demented (78.48% vs. 21.52%), and with a greater monthly decline of the ALSFRS-R (2.15 ± 2.87 points/months vs. 1.00 ± 1.36).

The factors influencing survival in these patients were PDC (HR 0.95, 95% C.I. 0.94-0.97, $p=0.001$), and CCI (HR 1.71, 95% C.I. 1.08-2.69, $p=0.021$).

The same results were obtained removing from the initial population patients who survived less than 4 months (46 patients) and less than 5 months (64 patients)(data not shown).

We investigated if treatment with riluzole and its duration were ineffective in the patients with the lowest ALSFRS-R score at diagnosis i.e. the lowest quartile according to this variable (ALSFRS-R global score < 35). In this group, multivariable analysis showed that factors predicting survival were PDC (HR 0.98, 95% C.I. 0.97-0.99, $p < 0.001$), weight loss at diagnosis (HR 1.07, 95% C.I. 1.03-1.11, $p < 0.001$), age at onset I (HR 1.05, 95% C.I. 1.02-1.08, $p = 0.002$), and diagnostic delay (HR 0.97, 95% C.I. 0.94-0.99, $p = 0.009$). When we limited the analysis to 25% of patients having the highest ALSFRS-R monthly decline (> 1.3 points/month), factors predicting survival were age at onset (HR 1.04, 95% C.I. 1.01-1.06, $p = 0.007$), dementia (HR 3.47, 95% C.I. 1.63-7.35, $p = 0.015$), weight loss at diagnosis (HR 1.06, 95% C.I. 1.01-1.10, $p = 0.009$), and gastrostomy (HR 2.47, 95% C.I. 1.35-4.50, $p = 0.003$).

DISCUSSION

Riluzole is the only drug approved for the treatment of ALS by the European Medical Agency in Europe. After >20 years from its discovery [28], a number of concerns about its therapeutic effects persist: the relatively unknown mechanism of action, the modest prolongation of survival (on average a few months), with concerns about cost effectiveness, the lack of benefit on some secondary measures of efficacy, and some controversies related to the drug efficacy in population-based studies and clinical practice [11]. Otherwise ALS has no cure, currently, and riluzole has a satisfactory safety profile, leading to recommend the drug to slow disease progression for patients with ALS [29].

Because ALS treatment (including all therapeutic interventions and drugs) is completely covered by INHS and considering that it has a relatively low cost, the percentage of ERR residents with ALS using riluzole is very high. In our study, 84% of patients received riluzole treatment for at least one month, a fraction that is higher than that reported in other studies carried on in Europe [30], USA [31] and Asia [17] probably due to the different socio-economic context. Given the widespread use of this drug in our country it is therefore important to determine whether and to what extent it is effective in the real-world ALS population, to improve patient counseling and the design of clinical trials.

In previous non-RCTs studies, the effect of riluzole on ALS survival was controversial, with some studies showing no gain of survival [32], and others showing an effect ranging from 6 to 12 months [3, 5, 12], a few studies documenting an effect for the first 6 months of treatment only, with a 15% reduction in mortality at 6 months [16]. Some investigators suggested that riluzole could only slow down motor neuron degeneration and it is more effective in early-

1 stage patients [15, 33]; others suggest a possible beneficial effect only after long-term use of the drug [17]. This
2 different effectiveness may be attributed to the different target populations (registry populations or referral cohorts).
3
4 Moreover, in most studies riluzole use was analyzed as a categorical factor (use versus non use)[3, 5, 12, 15, 34–37],
5
6 whereas only two recent studies considered the cumulative defined daily dose [16, 17] coming to opposite conclusions:
7
8 a limited effect for the first 6 months in Austrian ALS patients [16], a possible effect due to long-term use in a Chinese
9
10 cohort [17].

11 **In our study, analysing survival in dependence of riluzole treatment showed a beneficial effect of the drug for the initial**
12 **months of therapy only, because at 24 months after diagnosis the survival curves of riluzole-treated and untreated**
13 **patients crossed with untreated patients showing an apparently better survival thereafter (figure 1A), as already**
14 **reported [12, 16]. Although patients who took riluzole did not survive significantly more than patients who did not,**
15 **prognosis of patients who took riluzole for more time (in relation to disease duration) was better. Our apparently**
16 **inconsistent results, in agreement with a recent study [17], may reflect the hypothesis that drug efficacy can be**
17 **confirmed only after long-term treatment.** Moreover, patients enrolled in riluzole RCTs were perhaps more motivated to
18
19 carry on the study treatment until the study end (hoping in a new therapy for ALS) than current real-world patients who
20
21 may discontinue the drug because they saw themselves worsening or were aware of its limited effectiveness [37].
22
23 Nevertheless, since the duration of treatment with riluzole would be longer in patients living longer, this might be a
24
25 potential confounder of the dose-dependent effect of riluzole on survival, and the lack of difference between patients
26
27 treated with riluzole and patients who were untreated may be due to the fact that more severely affected individuals do
28
29 not take it or discontinue its intake because of a rapidly worsening condition. For this reason, we also adjusted our
30
31 analysis for immortal time bias, but results were similar, suggesting that this potential bias did not substantially affect
32
33 our findings.
34
35

36
37 We also eliminated from our sample patients died within 6 months from diagnosis, but the results were unchanged.
38
39 To test the hypothesis of a lack of efficacy of riluzole treatment in more severely affected patients at diagnosis, we also
40
41 evaluated patients with a low ALSFRS-R score at diagnosis: in that group, the extent of treatment duration
42
43 independently influenced survival.
44
45

46
47 **Conversely, our analyses in patients who had a high ALSFRS-R monthly decline showed that in this small sample PDC**
48 **did not influenced survival: in this rapidly progressive population riluzole, whenever given, has no effect on survival.**
49

50
51 As efficacy of riluzole seemed related to the duration of its intake, we hypothesized that the earlier it is given to patients
52
53 the longer the patients will survive, as already suggested [38]. Therefore we studied the relationship between delay from
54
55 disease onset and start of riluzole, and survival. We found that the earlier the patients took riluzole, the worse the
56
57 prognosis, probably because of the well known negative prognostic role of a short diagnostic delay in ALS [3].
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

On the contrary, we identified the characteristics of patients who took riluzole for a longer period of time: these patients were younger, more frequently with spinal onset, without cognitive impairment, and with a lower CCI. Younger age and spinal onset, and absence of dementia, are well known factors related to a better survival [3]. These patients may be more compliant and more prone to comply with medical counseling and pharmacological treatment: we found that patients treated for a longer time with riluzole underwent more frequently NIV and IV, suggesting a general attitude to accept all therapeutic interventions which may influence survival [3]. These findings are similar to what reported by others [17].

The study has strengths and limitations. The major strength is its population-based design. In addition, we are fairly confident that case ascertainment in the study area was almost complete, as shown by the incidence rate of the disease [18].

A limitation of this study is the observational design, compared to the experimental context of RCTs. However, observational studies particularly if population-based, have the advantage of a longer follow-up than the RCTs, and also include participants who approximate real world population [39]. This also means that, since age at onset and disease severity are more heterogeneous in a population-based registry study than in RCTs [28, 40, 41], older and more severely affected patients were included in our study in comparison with RCTs. A recent Cochrane review [11] showed that analyzing pooled riluzole RCTs data, the effects of riluzole on survival were significant when the homogeneous group of participants in the first two trials were considered [28, 40], but when all three trials were analyzed [41], there was a high heterogeneity due to the addition of more seriously affected and older patients, and the combined treatment effect fell just short of significance [11].

Another possible limitation of the study is the fact that the vast majority of ALS patients received Riluzole [23]. In addition, we did not collect data on advance directives of patients in relation to nutritional and respiratory supports which can influence survival and we do not have data about reasons for riluzole treatment discontinuations or related side effects. Furthermore, we had genetic status only in a limited number of cases and this information would have impact survival especially in relation to C9orf72 hexanucleotide repeat expansion [42].

Finally, as discussed before, a specific and important weakness of our study is that, since the duration of treatment would be greater in patients living longer, this may be a potential confounder of the dose-dependent effect of riluzole on survival.

In summary, in our study patients who received riluzole for a proportionally longer period of time survived longer, thus suggesting that long-term use of riluzole may prolong survival. Younger and cognitively normal ALS patients and patients with a spinal onset, a lower CCI, and lower decline of ALSFRS-R were more likely to use riluzole for a longer time and survived longer. However, further studies are needed to verify if long-term use of riluzole prolongs survival.

REFERENCES

1. Chiò A, Calvo A, Moglia C, Mazzini L, Mora G, PARALS study group. (2011) Phenotypic heterogeneity of amyotrophic lateral sclerosis: a population based study. *J Neurol Neurosurg Psychiatry* 82:740–6.
2. Al-Chalabi A, Hardiman O, Kiernan MC, et al. (2016) Amyotrophic lateral sclerosis: moving towards a new classification system. *Lancet Neurol* 15:1182–1194. doi: 10.1016/S1474-4422(16)30199-5
3. Calvo A, Moglia C, Lunetta C, et al. (2017) Factors predicting survival in ALS: a multicenter Italian study. *J Neurol* 264:54–63. doi: 10.1007/s00415-016-8313-y
4. Rooney J, Byrne S, Heverin M, Tobin K, Dick A, Donaghy C, Hardiman O (2015) A multidisciplinary clinic approach improves survival in ALS: a comparative study of ALS in Ireland and Northern Ireland. *J Neurol Neurosurg Psychiatry* 86:406–501.
5. Georgouloupoulou E, Fini N, Vinceti M, et al. (2013) The impact of clinical factors, riluzole and therapeutic interventions on ALS survival: A population based study in Modena, Italy. *Amyotroph Lateral Scler Frontotemporal Degener* 8421:1–8. doi: 10.3109/21678421.2013.763281
6. Mitsumoto H, Brooks BR, Silani V (2014) Clinical trials in amyotrophic lateral sclerosis: Why so many negative trials and how can trials be improved? *Lancet Neurol* 13:1127–1138. doi: 10.1016/S1474-4422(14)70129-2
7. Hardiman O, van den Berg LH, Kiernan MC (2011) (2011) Clinical diagnosis and management of amyotrophic lateral sclerosis. *Nat Rev Neurol* 7:639–649.
8. Vucic S, Lin CS, Cheah BC, Murray J, Menon P, Krishnan AV KM (2013) Riluzole exerts central and peripheral modulating effects in amyotrophic lateral sclerosis. *Brain* 136:1361–70.
9. Gutierrez J, Federici T, Peterson B, Bartus R, Betourne A, Boulis NM (2016) Development of Intrathecal Riluzole: A New Route of Administration for the Treatment of Amyotrophic Lateral Sclerosis Patients. *Neurosurgery* 63:193.
10. Dyer AM (2017) Riluzole 5 mg / mL oral suspension : for optimized drug delivery in amyotrophic lateral sclerosis. 59–64. doi: 10.2147/DDDT.S123776
11. Miller RG, Mitchell JD, Moore DH (2012) Riluzole for amyotrophic lateral sclerosis (ALS)/motor neuron disease (MND)(Review). *Forbes* 3:1–34. doi: CD001447 [pii]r10.1002/14651858.CD001447
12. Zoccolella S, Beghi E, Palagano G, et al. (2007) Riluzole and amyotrophic lateral sclerosis survival: A population-based study in southern Italy. *Eur J Neurol* 14:262–268. doi: 10.1111/j.1468-1331.2006.01575.x
13. Mitchell JD, O'Brien MR, Joshi M (2006) Audit of outcomes in motor neuron disease (MND) patients treated with riluzole. *Amyotroph Lateral Scler Other Mot Neuron Disord* 7:67–71.

14. Turner MR, Bakker M, Sham P, Shaw CE, Leigh PN, Al-Chalabi A (2001) The Kings' Database 1999 – 2000: an analysis of the effect on survival of interventions in amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Mot Neuron Disord* 2:43.
15. Traynor BJ, Alexander M, Corr B, Frost E, Hardiman O (2003) An outcome study of riluzole in amyotrophic lateral sclerosis--a population-based study in Ireland, 1996-2000. *J Neurol* 250:473–479.
16. Cetin H, Rath J, Füzi J, et al. (2015) Epidemiology of amyotrophic lateral sclerosis and effect of riluzole on disease course. *Neuroepidemiology* 44:6–15. doi: 10.1159/000369813
17. Chen L, Liu X, Tang L, et al. (2016) Long-Term Use of Riluzole Could Improve the Prognosis of Sporadic Amyotrophic Lateral Sclerosis Patients : A Real-World Cohort Study in China. 8:1–8. doi: 10.3389/fnagi.2016.00246
18. Mandrioli J, Biguzzi S, Guidi C, et al. (2014) Epidemiology of amyotrophic lateral sclerosis in Emilia Romagna Region (Italy): A population based study. 8421:262–268. doi: 10.3109/21678421.2013.865752
19. Brooks BR, Miller RG, Swash M, Munsat TL (2000) El Escorial revisited: revised criteria for the diagnosis of amyotrophic lateral sclerosis. *Amyotroph Lateral Scler Other Motor Neuron Disord* 1:293–299. doi: DOI 10.1080/146608200300079536
20. Franchignoni F, Mora G, Giordano A, et al. (2013) Evidence of multidimensionality in the ALSFRS-R Scale: a critical appraisal on its measurement properties using Rasch analysis. *J Neurol Neurosurg Psychiatry* 84:1340–1345. doi: 10.1136/jnnp-2012-304701
21. Franchignoni F, Mandrioli J, Giordano A, Ferro S (2015) A further Rasch study confirms that ALSFRS-R does not conform to fundamental measurement requirements. *Amyotroph Lateral Scler Front Degener* 8421:1–7. doi: 10.3109/21678421.2015.1026829
22. Fasano A, Fini N, Ferraro D, et al. (2017) Percutaneous endoscopic gastrostomy, body weight loss and survival in amyotrophic lateral sclerosis: a population-based registry study. *Amyotroph Lateral Scler Front Degener* 0:1–10. doi: 10.1080/21678421.2016.1270325
23. Andersen PM, Abrahams S, Borasio GD, et al. (2012) EFNS guidelines on the Clinical Management of Amyotrophic Lateral Sclerosis (MALS) - revised report of an EFNS task force. *Eur J Neurol* 19:360–375. doi: 10.1111/j.1468-1331.2011.03501.x
24. Fini N, Georgouloupoulou E, Vinceti M, et al. (2014) Noninvasive and invasive ventilation and enteral nutrition for ALS in Italy. *Muscle Nerve* 50:508–516.
25. Suissa S (2008) Immortal time bias in pharmaco-epidemiology. *Am J Epidemiol* 167:492-9.
26. Labra J, Menon P, Byth K, et al. (2015) Rate of disease progression: a prognostic biomarker in ALS. *J Neurol*

Neurosurg Psychiatry jnnp-2015-310998. doi: 10.1136/jnnp-2015-310998

- 1
2 27. Thygesen SK, Christiansen CF, Christensen S, Lash TL, Sørensen HT (2011) The predictive value of ICD-10
3 diagnostic coding used to assess Charlson comorbidity index conditions in the population-based Danish
4 National Registry of Patients. *BMC Med Res Methodol* 11:83. doi: 10.1186/1471-2288-11-83.
5
- 6
7 28. Bensimon G, Lacomblez L, Meininger V. (1994) A controlled trial of riluzole in amyotrophic lateral sclerosis.
8 *ALS/Riluzole Study Group*. 330:585–91.
9
- 10
11 29. Miller RG, Jackson CE, Kasarskis EJ, et al. (2009) Practice parameter update: the care of the patient with
12 amyotrophic lateral sclerosis: multidisciplinary care, symptom management, and cognitive/behavioral
13 impairment (an evidence-based review): report of the Quality Standards Subcommittee of the American
14 Neurology 73:1227–33.
15
16
17
18
19 30. Cetin H, Klickovic U, Rath J, et al. (2015) Associations between co-medications and survival in ALS — a
20 cohort study from Austria. 1698–1705. doi: 10.1007/s00415-015-7767-7
21
22
23 31. Gordon PH (2013) Amyotrophic Lateral Sclerosis: An update for 2013 Clinical Features, Pathophysiology,
24 Management and Therapeutic Trials. *Aging Dis* 4:295–310. doi: 10.14336/AD.2013.0400295
25
26
27 32. Sívori M, Rodríguez GE, Pascansky D, Sáenz C, Sica RE (2007) Outcome of sporadic amyotrophic lateral
28 sclerosis treated with non-invasive ventilation and riluzole. *Med (B Aires)* 67:326–330.
29
30
31 33. Tavakoli M (2002) Disease progression in amyotrophic lateral sclerosis. Identifying the cost-utility of riluzole
32 by disease stage. *Eur J Heal Econ* 3:156–165.
33
34
35 34. Chio A, Logroscino G, Hardiman O, et al. (2009) Prognostic factors in ALS: A critical review. *Amyotroph*
36 *Lateral Scler* 10:310–323. doi: 10.3109/17482960802566824
37
38
39 35. Chio A, Mora G, Leone M, et al. (2002) Early symptom progression rate is related to ALS outcome: a
40 prospective population-based study. *Neurology* 59:99–103. doi: 10.1212/WNL.59.12.2012-a
41
42
43 36. Kiernan MC, Vucic S, Cheah BC, et al. (2011) Amyotrophic lateral sclerosis. *Lancet* 377:942–955. doi:
44 10.1016/S0140-6736(10)61156-7
45
46
47 37. Lee CT-C, Chiu Y-W, Wang K-C, et al. (2013) Riluzole and Prognostic Factors in Amyotrophic Lateral
48 Sclerosis Long-term and Short-term Survival: A Population-Based Study of 1149 Cases in Taiwan. *J Epidemiol*
49 23:35–40. doi: 10.2188/jea.JE20120119
50
51
52 38. Knibb JA, Keren N, Kulka A, et al. (2016) A clinical tool for predicting survival in ALS. *J Neurol Neurosurg*
53 *Psychiatry* 87:1361–1367. doi: 10.1136/jnnp-2015-312908
54
55
56 39. Chiò A, Canosa A, Gallo S, et al. (2011) ALS clinical trials: do enrolled patients accurately represent the ALS
57 population? *Neurology* 77:1432–1437. doi: 10.1212/WNL.0b013e318232ab9b
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
40. Lacomblez L, Bensimon G, Leigh PN, Guillet P, Meininger V (1996) Dose-ranging study of riluzole in amyotrophic lateral sclerosis. *Amyotrophic Lateral Sclerosis/Riluzole Study Group II. Lancet* 347:1425–1431.
41. Bensimon G, Lacomblez L, Delumeau JC, Bejuit R, Truffinet P, Meininger V; Riluzole/ALS Study Group II.. (2002) A study of riluzole in the treatment of advanced stage or elderly patients with amyotrophic lateral sclerosis. *J Neurol* 249:609–615.
42. Rooney J, Fogh I, Westeneng HJ, et al. (2017) C9orf72 expansion differentially affects males with spinal onset amyotrophic lateral sclerosis. *J Neurol Neurosurg Psychiatry* 88:281.

TABLES AND FIGURES LEGENDS

Table 1: Patients characteristics (681 patients)

Table 2: Clinical factors and survival (from onset/diagnosis to death or tracheostomy) in incident patients of Emilia Romagna, Italy (univariate analysis) (681 patients)

Table 3: Clinical characteristics of patients with respect to % of riluzole treatment duration in relation to survival ($\geq 90\%$ versus 1-90% of time of disease duration, versus no treatment) (681 patients)

Table 4: Independent prognostic factors (multivariable Cox analysis)

Figure 1: A) Tracheostomy-free survival from diagnosis of incident ALS cases in Emilia Romagna Region based on riluzole treatment; B) Tracheostomy-free survival from diagnosis of incident ALS cases in Emilia Romagna Region based on percentage of time of riluzole treatment in relation to disease duration ($\geq 75\%$ versus $< 75\%$ of time of disease duration); C) Tracheostomy-free survival from diagnosis of incident ALS cases in Emilia Romagna Region based on percentage of time of riluzole treatment in relation to disease duration ($\geq 90\%$ versus $< 90\%$ of time of disease duration).

Figure 1




Table 1: Patients' characteristics (N = 681)

<i>Explanatory variables</i>	Total N=681 n (%) m [SD]^a	Men N=371 n (%) m [SD]	Women N=310 n (%) m [SD]	p-value
ALS Onset (Bulbar)	238 (34.95)	107 (28.84)	131 (42.26)	<0.001
Age at onset	66.50 [±11.09]	65.87 [±10.66]	67.26 [±11.57]	0.102
Diagnostic delay	13.16 [±13.09]	13.13 [±13.85]	13.20 [±12.14]	0.947
Phenotype				
Bulbar	211 (30.98)	92 (24.80)	119 (38.39)	<0.001
Classic	286 (42.00)	177 (47.71)	109 (35.16)	
Flail arm and leg	119 (17.47)	68 (18.33)	51 (16.45)	
UMNp ^c	38 (5.58)	16 (4.31)	22 (7.10)	
Respiratory	21 (3.96)	15 (4.04)	6 (1.94)	
Unknown	6 (0.88)	3 (0.81)	3 (0.97)	
Revised El Escorial criteria				
Definite	189 (27.75)	94 (25.34)	95(30.65)	0.250
Clinically probable	216 (31.72)	122 (32.88)	94 (30.32)	
Probable lab-supported	92 (13.51)	48 (12.94)	44 (14.19)	
Possible	136 (19.97)	82 (22.10)	54 (17.42)	
Dementia	64 (9.39)	31 (8.36)	33 (10.65)	0.308
BMI^d at diagnosis (Kg/m²)	24.39 [±3.98]	24.94 [±3.72]	23.73 [±4.19]	<0.001
Riluzole (Yes)	573 (84.14)	319 (85.98)	254 (81.94)	0.150
Treatment duration (days)	869.24 [±645.00]	867.84 [±639.67]	870.99 [±652.84]	0.954
Delay from onset to riluzole intake (days)	421.13 [±372.12]	408.82 [±367.45]	436.48 [±378.02]	0.378
Delay from diagnosis to riluzole intake (days)	41.53 [±97.98]	39.74 [±80.31]	43.76 [±116.46]	0.626
Gastrostomy (Yes)	207 (30.39)	96 (25.88)	111 (35.81)	0.005

Non invasive ventilation (Yes)	246 (36.12)	147 (39.62)	99 (31.94)	0.038
Invasive ventilation (Yes)	103 (15.15)	65 (17.52)	38 (12.26)	0.059
ALSFRS-R^e at diagnosis	38.58 [\pm 7.95]	39.28 [\pm 7.26]	37.75 [\pm 8.63]	0.013
ALSFRS-R monthly decline (points/month) measured at diagnosis	1.14 [\pm 1.66]	1.08 [\pm 1.41]	1.21 [\pm 1.92]	0.300
Charlson Comorbidity Index	2.73 [\pm 1.56]	2.69 [\pm 1.50]	2.77 [\pm 1.62]	0.221

SD^a: Standard Deviation; NA^b: Not Appropriate; UMN-P^c: Upper Motor Neuron predominant; BMI^d: Body Mass

Index; ALSFRS-R^e: ALS Functional Rating Scale – Revised.

Table 2: Demographic and clinical factors and survival (from onset/diagnosis to death or tracheostomy) in incident patients of Emilia Romagna, Italy (univariate analysis) (N = 681)

Variable	Categories	Survival from onset			Survival from diagnosis		
		HR ^c	95%CI ^d	p-value	HR	95%CI	p-value
Sex	Woman	1	(reference)	0.239	1	(reference)	0.308
	Man	1.12	0.93-1.37		1.11	0.91-1.35	
Onset	Spinal	1	(reference)	<0.001	1	(reference)	<0.001
	Bulbar	1.77	1.45-2.16		1.69	1.39-2.06	
Diagnostic delay	Months	0.41	0.33-0.50	<0.001	0.98	0.97-0.99	<0.001
Age at onset	Years	1.04	1.03-1.05	<0.001	1.04	1.03-1.05	<0.001
Phenotype	Bulbar	1	(reference)	<0.001	1	(reference)	0.015
	Classic	0.63	0.50-0.78		0.80	0.64-1.00	
	Flail arm/leg	0.47	0.35-0.64		0.70	0.52-0.94	
	Upper MN-predominant	0.32	0.19-0.56		0.52	0.31-0.86	
	Respiratory	1.29	0.77-2.16		1.15	0.65-2.09	
Dementia	No	1	(reference)	<0.001	1	(reference)	<0.001
	Yes	2.02	1.50-2.73		2.26	1.67-3.07	
Familial ALS	No	1	(reference)	0.711	1	(reference)	0.679
	Yes	1.09	0.68-1.75		1.10	0.69-1.771	
EEC-R ^a	Definite	1	(reference)	<0.001	1	(reference)	0.025
	Probable	0.71	0.56-0.91		0.89	0.70-1.13	
	Probable lab. supported	0.43	0.31-0.61		0.68	0.49-0.95	
	Possible	0.62	0.47-0.82		0.68	0.51-0.92	
ALSFRS-R ^b at diagnosis	Points	0.99	0.97-1.00	0.032	0.98	0.97-0.99	<0.001
ALSFRS-R monthly decline	Points	1.32	1.27-1.38	<0.001	1.24	1.10-1.39	<0.001
ALSFRS-R monthly decline	<1 point/month	1	(reference)	<0.001	1	(reference)	0.014
	≥1 point/month	3.75	2.97-4.74		1.33	1.06-1.65	

BMI at diagnosis	Kg/m ²	0.98	0.95-1.01	0.118	0.97	0.95-1.00	0.086
Weight variation between healthy state and diagnosis	kg	1.04	1.03-1.05	<0.001	1.05	1.04-1.07	<0.001
Riluzole treatment (No/Yes)	No	1	(reference)	0.39	1	(reference)	0.703
	Yes	1.12	0.86-1.48		1.05	0.80-1.39	
Riluzole treatment duration/disease duration (onset to death/tracheostomy)	<60%	1	(reference)	<0.001	NA	NA	NA
	≥60%	0.57	0.47-0.69		NA	NA	NA
Riluzole treatment duration/disease duration (onset to death/tracheostomy)	%	0.49	0.37-0.66	<0.001	NA	NA	NA
Riluzole treatment duration/disease duration (diagnosis to death/tracheostomy)	<75%	NA	NA	NA	1	(reference)	0.001
	≥75%	NA	NA	NA	0.69	0.55-0.86	
Riluzole treatment duration/disease duration (diagnosis to death/tracheostomy)	%	NA	NA	NA	0.99	0.99-1.00	0.010
Delay from onset to riluzole intake	months	0.96	0.95-0.97	<0.001	0.98	0.97-0.99	<0.001
Delay from onset to riluzole intake	<10 months	1	(reference)	<0.001	1	(reference)	<0.001
	≥10 months	0.43	0.35-0.53		0.64	0.52-0.79	
Non invasive ventilation	No	1	(reference)	<0.001	1	(reference)	<0.001
	Yes	1.58	1.30-1.92		1.59	1.318-1.94	
Gastrostomy	No	1	(reference)	<0.001	1	(reference)	<0.001
	Yes	1.86	1.53-2.27		1.80	1.47-2.19	
Charlson Comorbidity Index	Points	1.21	1.15-1.29	<0.001	1.24	1.17-1.31	<0.001

^aEEC-R: El Escorial Criteria- Revised; ^bALSFRS-R: ALS Functional Rating Scale –Revised; ^cHR= Hazard Ratio; ^dCI=

Confidence Interval; Significant results in bold

Table 3: Clinical characteristics of patients with respect to % of riluzole treatment duration in relation to survival from diagnosis (>90% versus 1-90% of time of disease duration, versus no treatment) (N = 681)

<i>Explanatory variables</i>	No riluzole treatment N=106 n (%) m [SD] ^d	Riluzole treatment ≤90% N=228 n (%) m [SD]	Riluzole treatment >90% N=347 n (%) m [SD]	p-value
ALS Onset (Bulbar)	44 (41.51)	88 (38.59)	106 (30.55)	0.043
Sex (Man)	51 (48.11)	125 (54.82)	195 (56.19)	0.340
Age at onset	69.85 [±11.04]	68.33 [±9.99]	64.27 [±11.35]	<0.001
Diagnostic delay	15.77 [±17.98]	12.34 [±13.30]	12.81 [±10.85]	0.067
Phenotype				
Bulbar	42 (39.62)	73 (32.02)	97 (27.95)	0.187
Classic	41 (38.68)	100 (43.86)	150 (43.23)	
Flail arm and leg	14 (13.21)	32 (14.03)	73 (21.04)	
UMNp ^a	6 (5.66)	13 (5.70)	19 (5.47)	
Respiratory	3 (2.83)	10 (4.38)	8 (2.30)	
Revised El Escorial Criteria				
Definite	27 (28.72)	71 (33.65)	91 (27.74)	0.562
Clinically probable	37 (39.36)	71 (33.65)	108 (32.93)	
Probable lab-supported	12 (12.76)	26 (12.32)	54 (16.46)	
Possible	18 (19.15)	43 (20.38)	75 (22.86)	
Dementia	13 (12.26)	32 (14.03)	19 (5.47)	0.001
BMI^b at diagnosis (Kg/m²)	24.00 [±4.71]	24.16 [±4.04]	24.64 [±3.75]	0.290
Gastrostomy (Yes)	24 (22.64)	78 (34.21)	105 (30.26)	0.101
Non invasive ventilation (Yes)	28 (26.41)	77 (33.77)	141(40.63)	0.019
Invasive ventilation (Yes)	8 (7.55)	49 (21.58)	46 (13.26)	0.001
ALSF^c monthly decline (points/month)	0.83 [±0.69]	0.86 [±0.85]	0.69 [±0.67]	0.016
Days from diagnosis to riluzole intake	NA	78.98 [±148.67]	17.62 [±16.04]	<0.001
Days from onset to riluzole intake	NA	442.37 [±438.73]	401.91 [±326.05]	<0.001
Absolute duration of Riluzole treatment	NA	361.38 [±368.55]	1027.25 [±	<0.001

(days)			611.16]	
Charlson Comorbidity Index				0.027
<3	41 (38.68)	88 (38.60)	173 (49.86)	
3-5	59 (55.66)	127 (55.70)	165 (47.55)	
6-10	6 (5.66)	13 (5.70)	9 (2.59)	

^aUMN-p= Upper Motor Neuron- predominant, ^bBMI= Body Mass Index, ^cALSFRR-R: ALS Functional Rating Scale –

Revised; ^dSD= Standard Deviation; Significant results in bold

Table 4: Independent prognostic factors (multivariable Cox analysis)

Variable	Survival from diagnosis to death or tracheostomy		
	Hazard Ratio	95%CI	p-value
Age at onset (years)	1.04	1.02-1.05	<0.001
Dementia (yes/no)	1.56	1.05-2.32	0.027
Revised El Escorial Criteria classification	0.88	0.78-0.98	0.021
Weight variation between healthy state and diagnosis (kg)	1.05	1.03-1.07	<0.001
Gastrostomy (Yes/No)	1.46	1.14-1.88	0.003
Non invasive ventilation (Yes/No)	1.43	1.12-1.82	0.004
Time from onset to diagnosis (months)	0.98	0.97-0.99	0.004
Riluzole treatment duration/disease duration (diagnosis to death/tracheostomy) (%)	0.98	0.98-0.99	<0.001

^aALSFRS-R: ALS Functional Rating Scale - Revised. Significant results in bold